



# Heart Rate Variability as a Non-Invasive Method to Assess Welfare in Dairy Cows



Elizabeth Regina Carvalho\* and Guilherme Galhardo Franco

College of Veterinary Medicine and Agricultural Sciences, São Paulo State University (UNESP), Brazil

\*Corresponding author: Elizabeth Regina Carvalho, College of Veterinary Medicine and Agricultural Sciences, Jaboticabal, Brazil, Email: [beth\\_rcarvalho@hotmail.com](mailto:beth_rcarvalho@hotmail.com)

Submission: 📅 February 03, 2018; Published: 📅 March 08, 2018

## Abstract

Adaptation to environmental changes can be challenging for highly productive dairy cow breeds. Animal welfare associated with husbandry procedures can be assessed not only by classical descriptive behavioral observations, but with physiological measures as well. Assessment of blood cortisol levels requires handling, what represents additional stress to cows. Measures of cardiovascular parameters including heart rate (HR) and heart rate variability (HRV) have a long tradition as indicators of health and welfare in livestock species since the beginning of the 1970s. HRV is a parameter that reflects the balance between sympathetic and parasympathetic nervous activity. Therefore, the power spectral analysis of HRV allows researchers to measure stress levels in cattle without handling or restraining them, thus making it a non-invasive indicator of welfare.

**Keywords:** Autonomic nervous system; Cattle; ECG; Stress; Sympathetic; Parasympathetic

**Abbreviations:** ANS: Autonomic Nervous System; ECG: Electrocardiogram; HF: High Frequency; HPA: Hypothalamic-Pituitary-Adrenal Axis; HR: Heart Rate; HRV: Heart Rate Variability; LF: Low Frequency.

## Introduction

Adaptation to environmental changes can be challenging for highly productive dairy cow breeds, once intensive dairy farming, housing and milking systems are main factors in determining the welfare of animals [1]. The impact of technological environment on cattle welfare has been examined in many different contexts. Certain welfare studies showed that for intensively farmed dairy cows, the fear due to routine handling [2], milking technology [3,4] and painful situations [5,6] mean a stressful load [7], which consequently have negative impacts on milk production [3]. Animal welfare associated with husbandry procedures can be assessed not only by classical descriptive behavioral observations [8,9], but with physiological measures as well [10-12].

Measures of cardiovascular parameters including heart rate (HR) and heart rate variability (HRV) have a long tradition as indicators of health and welfare in livestock species since the beginning of the 1970s. In farm animals, the vagal component of the autonomic nervous system (ANS) plays a key role in regulating HR in response to stress [2,12]. Many parameters of HRV give information about cardiac vagal tone and the sympathetic-parasympathetic balance [3]. Consequently, besides traditional ways of assess stress – assay of cortisol in plasma, serum or feces [13], assessment of HR and HRV has also been investigated in dairy cattle in veterinary, behavioral and applied animal research worldwide [8-15].

## Significance of HR and HRV as Non-Invasive Stress Parameters in Dairy Cows

There are two major physiological pathways reported to be involved in stress responses in mammals. [1] Stress increases the activity of the hypothalamic-pituitary-adrenal (HPA) axis, which leads to an elevation in circulating levels of cortisol [16]. Thus, serum cortisol level is one of the parameters to evaluate activity of the HPA axis and therefore, is used to estimate stress intensity in farm animals [7,17]. Assessment of blood cortisol levels requires handling, what represents additional stress to cows. [2] The other physiological response is related to ANS: stress increases sympathetic nervous activity and decreases parasympathetic nervous tonus [16].

HR is defined as the number of heart beats per minute, and interpretations have often been based on the assumption that HR reflects the activity of the sympathetic branch of the ANS, and therefore as an indicator of the stress response [18,19]. However, increases on HR can occur either in a state of pleasure/happiness or in response to a negative/painful stimulus. The complex interplay of the two branches – sympathetic and parasympathetic – of the ANS is not always comprehensible when cardiac activity is measured only by HR [20,21], as rise in HR could be attributable to an increase in sympathetic activity, decrease of vagal tone or the simultaneous changes in both systems [19].

Heart rate variability is a parameter that reflects the balance between sympathetic and parasympathetic nervous activity [1, 22, 23]. In the power spectrum of HRV, the high frequency (HF) power corresponds to the respiratory frequency and is influenced by vagal activity [24] therefore HF power is an index of the parasympathetic nervous activity. The low frequency (LF) power is closely associated with the fluctuations of blood pressure [24], and is related to both sympathetic and parasympathetic nervous activity [25].

Previous studies have suggested that an analysis of HRV with a Holter-type electrocardiograph helps evaluate stress caused by sickness conditions in dairy cows [14, 26]. In calves with external stress caused by environment high temperature and insect harassment, or diarrhea, the HF power decreased and LF/HF ratio increased, indicating a reduction of parasympathetic nervous tone during stress load [25,27]. Analysis of HRV in another study [28] revealed that changes in the sympatho-vagal balance were clearly detected in bull calves following surgical castration with or without local anesthesia. Therefore, the power spectral analysis of HRV allows researchers to measure stress levels in cattle without handling or restraining them, thus making it a non-invasive indicator of welfare.

### Methods of Measurement and Analysis of HR and HRV in Dairy Cattle

Measuring HR and HRV is based on electrocardiography (ECG). Different types of Holter recorders, fixed or telemetric systems, as well as portable ECG monitors have been used to assess HR and HRV in dairy cows [1]. The latter ones were originally developed for human athletes and sport medicine research [23], and were found to be a valid and reliable method to HR measurement in animals [29].

In dairy cattle practice, the recording of interbeat intervals is accomplished with two specific transmitter electrodes, and a ECG monitor [1]. It is recommended to place one of the electrodes next to the sternum, on the left side of the chest (cardiac area) and the other one on the right scapula [1,23]. The contacting surface should be cleaned before attaching, and electrode belts can be easily fixed around the thorax with an elastic strap [1]. Signal receivers are usually fixed on the outside of girths. Another method is to fix receivers on the site of the observer, instead of the animals [1].

### Referances

- Kovács L, Jurkovich V, Bakony M, Szenci O, Póti P, et al. (2014 ) Welfare implication of measuring heart rate and heart rate variability in dairy cattle: literature review and conclusions for future research. *Animal* 8(2): 316-30.
- von Holst D (1998) The Concept of Stress and Its Relevance for Animal Behavior. *Adv Study Behav* 27: 1-131.
- Rushen J, Munksgaard L, Marnet PG, DePassillé AM (2001) Human contact and the effects of acute stress on cows at milking. *Appl Anim Behav Sci* 73(1): 1-14.
- Wenzel C, Schönreiter-Fischer S, Unshelm J (2003) Studies on step-kick behavior and stress of cows during milking in an automatic milking system. *Livest Prod Sci* 83(2-3): 237-46.
- Broom DM (1991) Animal welfare: concepts and measurement. *J Anim Sci* 69(10): 4167-75.
- Mellor DJ, Cook CJ, Stafford KJ (2000) Quantifying some responses to pain as a stressor. In: Mench GM (Ed.), *The biology of animal stress*. Wallingford, CABI Publishing, UK, pp. 171-198.
- Dantzer R, Mormède P(1983 ) Stress in farm animals: a need for reevaluation. *J Anim Sci* 57(1):6-18
- Millman ST (2013) Behavioral responses of cattle to pain and implications for diagnosis, management, and animal welfare. *Vet Clin North Am Food Anim Pract* 29(1):47-58
- Theurer ME, Amrine DE, White BJ (2013 )Remote noninvasive assessment of visceral pain associated with metritis in dairy cows. *J Anim Pract* 29(1):59-74.
- Stojkov J, von Keyserlingk MA, Marchant-Forde JN, Weary DM (2015) Assessment of visceral pain associated with metritis in dairy cows. *J Dairy Sci* 98(8): 5352-5361.
- Kovács L, Tózsér J, Szenci O, Póti P, Kézér FL, et al. (2014 ) Cardiac responses to palpation per rectum in lactating and nonlactating dairy cows. *J Dairy Sci* 97(11): 6955-6963.
- Hopster H, Blokhuis HJ (1994) Validation of a heart-rate monitor for measuring a stress response in dairy cows. *Can J Anim Sci* 74(3): 465-474.
- Jurkovich V, Kézér FL, Ruff F, Bakony M, Kulcsár M, et al. (2017) Heart rate, heart rate variability, faecal glucocorticoid metabolites and avoidance response of dairy cows before and after changeover to an automatic milking system. *Acta Vet Hung* 65(2): 301-313.
- Kovács L, Tózsér J, Bakony M, Jurkovich V (2013) Short communication: Changes in heart rate variability of dairy cows during conventional milking with nonvoluntary exit. *J Dairy Sci* 96(12):7743-7.
- Johns J, Patt A, Hillmann E. Do Bells (2015) Affect Behaviour and Heart Rate Variability in Grazing Dairy Cows? McElligott A (ed), 10(6): e0131632.
- Ulrich-Lai YM, Herman JP (2009) Neural regulation of endocrine and autonomic stress responses. *Nat Rev Neurosci* 10(6): 397-409.
- Mormède P, Andanson S, Aupérin B, Beerda B, Guémené D, et al. (2007) Exploration of the hypothalamic-pituitary-adrenal function as a tool to evaluate animal welfare. *Physiol Behav* 92(3): 317-339.
- Hopster H, O'Connell JM, Blokhuis HJ (1995) Acute effects of cow-calf separation on heart rate, plasma cortisol and behaviour in multiparous dairy cows. *Appl Anim Behav Sci* 44(1):1-8.
- Sgoifo A, Koolhaas JM, Musso E, De Boer SF (1999) Different Sympathovagal Modulation of Heart Rate During Social and Nonsocial Stress Episodes in Wild-Type Rats. *Physiol Behav* 67(5):733-738.
- Porges SW (1995) Cardiac vagal tone: a physiological index of stress. *Neurosci Biobehav Rev* 19(2): 225-233.
- Marchant-Forde RM, Marlin DJ, Marchant-Forde JN (2004) Validation of a cardiac monitor for measuring heart rate variability in adult female pigs: accuracy, artefacts and editing. *Physiol Behav* 80(4):449-58.
- Baselli G, Cerutti S, Civardi S, Liberati D, Lombardi F, et al. (1986) Spectral and cross-spectral analysis of heart rate and arterial blood pressure variability signals. *Comput Biomed Res* 19(6): 520-534.
- von Borell E, Langbein J, Després G, Hansen S, Lettieri C, et al. (2007) Heart rate variability as a measure of autonomic regulation of cardiac activity for assessing stress and welfare in farm animals - A review. *Physiol Behav* 92(3): 293-316.
- Akselrod S, Gordon D, Madwed JB, Snidman NC, Shannon DC, et al.(1985) Hemodynamic regulation: investigation by spectral analysis. *Am J Physiol* 249(4): H867-H875.


25. Bun C, Watanabe Y, Uenoyama Y, Inoue N, Ieda N, et al. (2018) Evaluation of heat stress response in crossbred dairy cows under tropical climate by analysis of heart rate variability. *J Vet Med Sci* 80(1):181-185.
26. Yoshida M, Onda K, Wada Y, Kuwahara M (2015) Influence of sickness condition on diurnal rhythms of heart rate and heart rate variability in cows. *J Vet Med Sci* 77(3):375-379.
27. Mohr E, Langbein J, Nürnberg G (2012) Heart rate variability: a noninvasive approach to measure stress in calves and cows. *Physiol Behav* 75(1-2): 251-259.
28. Stewart M, Verkerk GA, Stafford KJ, Schaefer AL, Webster JR (2010) Noninvasive assessment of autonomic activity for evaluation of pain in calves, using surgical castration as a model. *J Dairy Sci* 93(8): 3602-3609.
29. Essner A, Sjöström R, Ahlgren E, Lindmark B (2013) Validity and reliability of Polar® RS800CX heart rate monitor, measuring heart rate in dogs during standing position and at trot on a treadmill. *Physiol Behav* 114(115): 1-5.



Creative Commons Attribution 4.0  
International License

For possible submissions Click Here

[Submit Article](#)



### Approaches in Poultry, Dairy & Veterinary Sciences

#### Benefits of Publishing with us

- High-level peer review and editorial services
- Freely accessible online immediately upon publication
- Authors retain the copyright to their work
- Licensing it under a Creative Commons license
- Visibility through different online platforms