



Paraneoplastic Syndromes in Dogs

Almir Pereira de Souza* and Olívia Maria Moreira Borges

Veterinary Medicine Postgraduate Program, Brazil

*Corresponding author: Almir Pereira de Souza, Veterinary Medicine Postgraduate Program, Patos, PB, Brazil

Submission: February 12, 2018; Published: March 15, 2018

Abstract

Paraneoplastic Syndromes (PNS) are defined as a heterogeneous group of clinical and/or laboratory abnormalities that precede or occur simultaneously with a neoplasm or its metastases and can occur in both small animals and humans with comparatively similar symptoms. It is important to mention that PNS have already been described in the systems gastrointestinal, endocrine, circulatory/hematological, neurological, bone and nonspecific. Thus, recognizing and describing a PNS early on, whether common or not, is of great value in veterinary, given its importance in assisting the diagnosis and treatment of neoplastic processes.

Keywords: Neoplasm; Metastases; Diagnosis; Canine

Abbreviations: PNS: Paraneoplastic Syndromes

Introduction

Table 1: Occurrence of a few paraneoplastic syndromes in small animals and in humans.

Paraneoplastic Syndromes	Dog	Cat	Human
Myasthenia Gravis	Yes	Yes	Yes
Peripheral Neuropathy	Rare	Rare	Yes
Cerebellar degeneration	Yes	Yes	Yes
Hypoglycemia	Yes	Yes	Yes
Hypercalcemia	Yes	Yes	Yes
Anemia	Yes	Yes	Yes
Thrombocytopenia	Yes	Yes	Yes
Feline Paraneoplastic Alopecia	No	Yes	No
Nodular Dermatofibrosis	Yes	Yes	Yes
Acantose Nigricans	No	No	Yes
Sweet Syndrome	No	No	Yes
Cachexia	Yes	Yes	Yes
Hypertrophic Osteopathy	Yes	No	No

Paraneoplastic Syndromes (PNS) are defined as a heterogeneous group of clinical and/or laboratory abnormalities that precede or occur simultaneously with a neoplasm or its metastases. PNS can occur in both small animals and humans with comparatively similar symptoms (Table 1), being considered indirect side effects of cancer and may arise from humoral, hormonal, tumor or immune response to neoplasms, and characterized by ectopy of sites distant from the neoplastic site. Although important and described as important predictors or markers of possible regression or progression of oncologic disease, studies reporting the prevalence

of these alterations in veterinary are not available currently, being largely restricted to human medicine, where approximately 75% of cancer patients reportedly display paraneoplastic changes during the evolution of the neoplasia [1].

Therefore, PNS early recognition and clinical applicability are of great value in the clinical routine, since they may be the first sign of malignancy, and timely diagnosis can drastically influence patient prognosis and treatment [2]. Thus, the objective of this review is to describe the paraneoplastic syndromes in dogs, especially the less common PNS, since the knowledge and ability to recognize them early on can improve significantly the prognosis of cancer patients.

Paraneoplastic syndromes in dogs

PNSs are complex diseases/symptoms that may occur individually or in different body systems. To this end, it is important to mention that PNSs have already been described in the following systems: gastrointestinal (cachexia and gastrointestinal ulceration), endocrine (hypercalcemia, hypoglycemia and hyperestrogenism), circulatory/hematological (hypergammaglobulinemia, anemia, erythrocytosis, neutrophilic leukocytosis, leukopenia, disseminated intravascular coagulation (alopecia and nodular dermatofibrosis), neurological (myasthenia gravis and cerebellar degeneration), bone (peripheral osteopathy) and nonspecific (fever). In view of this clinical-hematological diversity, the formation mechanisms, clinical presentation and neoplastic origin of the less described and possibly frequent, but equally important paraneoplastic syndromes in the clinical routine of small animals are highlighted in the following paragraphs.

Hypertension and medullary hypoplasia forming mechanism

Excess estrogen interferes with the differentiation of hematopoietic stem cells, altering the iron use by erythrocyte precursors, and inhibiting production of erythrocyte stimulating factor in the circulation. Estrogenic marrow intoxication induces granulocytopenic increase and reduction of megakaryocytes and erythroid elements, leading initially to neutrophilia with left-sided deviation, thrombocytopenia and anemia, which may lead to aplasia or pancytopenia.

a. Neoplastic origin: Tumor of granulosa and supportive cells (sertolioma).

b. Clinical signs: Hematological, myelosuppression induced by hyperestrogenism, which is characterized by pancytopenia, may be observed. Clinically, the feminization syndrome shows slowly progressive alopecia, symmetrical, bilateral, and usually originating in the neck, lumbar region, perineum and genital region. Other findings include macular melanoma in the inguinal, perineal, and genital regions, and linear preputial dermatosis [3,4].

Nodular dermatofibrosis forming mechanism

Little known, but is believed to be of hereditary autosomal dominant origin and full penetrance.

a. Neoplastic origin: renal cystadenocarcinoma and uterine leiomyomas.

b. Clinical signs: Non-specific, but correlated to stages of uremia. The skin nodular lesions can be located in the trunk, head and distal portion of the limbs, especially between the digits and the cushions, when it leads to lameness [5].

Myasthenia gravis forming mechanism

The acquired form is triggered by the existence of antibodies against acetylcholine receptors in the neuromuscular junction, whereas the congenital form is the result of the hereditary deficiency of acetylcholine in the postsynaptic membranes of musculature.

a. Neoplastic origin: thymoma, osteosarcoma, lymphoma, biliary carcinoma and primary lung neoplasms.

b. Clinical signs: intermittent muscle weakness, exercise intolerance, dysphagia, megaesophagus, and respiratory paralysis [6].

Cerebellar degeneration: forming mechanism

Ectopic expression of neuronal antigens by the tumor tissue, triggering immune response against it, and cross-reaction against the antigens present in the nervous system, resulting in neurological manifestations.

a. Neoplastic origin: Lymphoma, mammary, pulmonary and ovarian carcinoma.

b. Clinical signs: ataxia and imbalances [7].

Thus, recognizing and describing a PNS early on, whether common or not, is of great value in veterinary, given its importance in assisting the diagnosis and treatment of neoplastic processes. Unusual paraneoplastic abnormalities such as hyperestrogenism, medullary hypoplasia, nodular dermatofibrosis, myasthenia gravis, and cerebellar degeneration should be remembered and considered in the evaluation of dogs with neoplasias since it could lead to a more adequate clinical management.

References


- Morrison WB (2002) Paraneoplastic syndromes and the tumors that cause them. In: Morrison WB (Ed.), *Cancer in dogs and cats*. (2nd edn), Williams e Wilkins, Baltimore, USA, pp. 731-743.
- Bergman PJ (2013) Paraneoplastic syndromes In: Withrow and MacEwen's (Eds.), *Small animal clinical oncology*. (5th edn), St. Saunders Elsevier, Louis, pp. 83-97.
- Bosschere HDE, Deprest C (2010) Estrogen-induced pancytopenia due to a Sertoli cell tumor in a cryptorchid Beauceron. *Vlaams Diergeneeskundig Tijdschrift* 79.
- Scott DW, Miller WH, Griffin CE (2001) In: Muller & Kirk's (Ed.), *Small animal dermatology*. (6th edn), Saunders, Philadelphia, USA.
- Langohr IM, Irigoyen LF, Salles, MWS, Kommers GD, Barros CSL (2002) Cistadeno carcinoma renal e dermatofibrose nodular em cães pastor alemão: 4 casos. *Cienc. Rural* 32(4): 621-626.
- Morrison WB (2007) Systemic Syndromes associated with cancer The North American Veterinary Conference, Orlando, Florida, USA.
- Bardy FB, Cagy M, Filho FP, lamarca JE, Rabinovitz M, et al. (2000) Degeneração cerebelar subaguda paraneoplásica - Relato de caso. *Arq neuropsiquiatr* 58(3A): 764-768.



Creative Commons Attribution 4.0 International License

For possible submissions Click Here

[Submit Article](#)



Approaches in Poultry, Dairy & Veterinary Sciences

Benefits of Publishing with us

- High-level peer review and editorial services
- Freely accessible online immediately upon publication
- Authors retain the copyright to their work
- Licensing it under a Creative Commons license
- Visibility through different online platforms