



IgY-Technology: A New Frontier in the Veterinary Diagnosing Procedures



Ferreira Junior A^{1*}, Santos JP², Santos MP², Sousa IO¹, Pereira FS² and Correia Lima AM²

¹Department of Animal Sanity and Production, University of Uberaba, Brazil

²Department of Veterinary Science, Federal University of Uberlândia, Brazil

*Corresponding author: Ferreira junior alvaro, Department of Animal Sanity and Production, University of Uberaba, Minas Gerais, Brazil, Email: alvaroferreirajr@gmail.com

Submission: 📅 December 23, 2017; Published: 📅 June 07, 2018

Abstract

Domestic animals, as bovine and equine, are vulnerable to diseases caused by infectious pathogens. Unfortunately, the pathogens that infect animals can also infect humans. In this context, the public health demands a global effort to control zoonotic diseases in herds from developing countries. Serology tests are applied to diagnose the infected animals, protect the human population and implement the disease control procedures. Among the reagents used in diagnostic platforms, the antibodies, as the Immunoglobulin G (IgG) and IgY from mammals and chickens, respectively, are some of them. The IgY antibodies are available in egg yolk content from immunized chicken. According to the principles of welfare, the large amount of antibodies and the antigenic target recognition made IgY an alternative biologic reagent in veterinary diagnosing tests. Innovative platforms for serology assays are being produced using IgY-technology.

Introduction

The hens' immune system has been a model for a myriad of revolutionary discoveries through the last decades. The bursa of Fabricius was the keystone for the research in humoral response's immunology. The absence of blood circulating antibodies after a bursectomy highlighted the lymphoid tissues function for B cells maturation and, also, their crucial role in antibody production. Additionally, hens' blood circulating antibodies were demonstrated to be transferred to their off spring by transport molecules on the surface of yolk membrane during the folliculogenesis.

The discovery that the antibody repertoire is the same in hens' blood and egg yolk brought an alternative to antibody production. Extracting antibodies from egg yolk of immunized hens reduces the painful events to the producing antibodies' host. Approaches were developed aiming the extraction of these antibodies by low cost and high concentration of antibodies taking in account animal welfare.

The hens' genome was sequenced and there were found coding genes for three classes of antibodies: Immunoglobulin (Ig) M (IgM), IgA and IgY. All of them are blood circulating antibodies. However, in the egg, a large number of IgY were found in yolk while only traces of IgM and IgA were deposited in white during egg formation. In this context, egg yolk was targeted for obtaining hens' antibodies. The IgY antibodies are progressively receiving attention from the scientific community, due to their similarities in biological functions with the mammals' IgG and their higher yield and easier extraction. In addition, when using avian antibodies in diagnostic approaches

the undesirable effects such as false positive results can be avoided. This feature is due to IgY structural difference in its Crystallizable Fragment (Fc) compared to the same equivalent region in IgG.

IgY is a promising biological molecule for many applications. Avian antibodies have been applied to diagnostic approaches such as viral and bacterial infections. Also, they have been used for immunotherapy such as oral passive immunization. IgY has been conjugated with enzymes, fluorophores or nano particles and they also can be used for enzyme-linked assays, immune histochemistry procedures or immunochromatography's test as well.

The bovine and equine herds are both susceptible to infectious diseases, including some zoonosis, which must be prevented by precocious diagnosing approaches. The detection of circulating or tissue antigens and specific blood circulating antibodies, brings celerity to diseases' diagnostic. In this context, a source for a quality biological molecule is needed. The IgY antibody fits all requirements to be this biological molecule. Hence, they have been used to diagnose viruses and bacteria that, infecting bovine and equine herds, cause risk to public health. In this mini review, we shall present recently published data using IgY-based approaches (named IgY-technology) for some challenges in veterinary medicine.

IgY antibodies: an overview

The hens' immune system is organized in lymphoid tissues, immune cells and molecules [1]. Bursa of Fabricius is an antibody-producing related lymphoid tissue; therefore, the bursectomy

drops the antibody production [2]. The B-cells derived plasma cells are the producing antibody cells. Firstly, compromised lymphoid stem cells living in bone marrow differentiate to immature B cells. The cell's maturation occurs in the Bursa of Fabricius tissue and it must involve a selective process preventing the development of autoimmune diseases [3]. The mature B cell leaves bursa of Fabricius to homing peripheral lymphoid tissues such as mucosa-associated lymphoid tissue being examples: eyes-associated Harderian's gland and gut-associated lymphoid tissue [4]. In these places, B cell must be exposed to foreign substances (named antigens), activate its genetic program for clonal expansion, and finally differentiate in Plasma cell [5]. B cell-derived Plasma cells are the antibodies' producing "factories".

The hens have three classes of antibodies: IgM, IgA and IgY, while mammals have five classes, IgM, IgA, IgG, IgE and IgD [6]. Avian IgY is an ancestor of IgG and IgE antibodies. The antibody molecule is a heterodimer protein assembled with two light chains and two heavy chains linked by inter chain disulfide bonds. Both, avian and mammalian primary immune response are featured by higher titles of IgM antibodies while the secondary immune response has mainly IgY (hens) or IgG (mammals) antibodies production [7,8]. Antibodies' coding genes control the hens and mammals' humoral immune response repertoire and the phylogenetic distance contributes to an interesting difference between these two animal groups [9].

Maternal antibodies are an important protection for offspring. Avian maternal antibodies are transferred to its descendent by egg yolk IgY while transplacental IgG transference occurs in mammals. During the hens' folliculogenesis there is an IgY translocation from the blood circulating to the egg yolk by specific receptors while IgM and IgA antibodies are secreted to the egg white [6,10].

Egg yolk antibodies are a promising source for high quality immunobiologics. The extraction of IgY antibodies from egg yolk reduces the number of painful events to the animal host. In addition, annually, just one hen can produce the equivalent number of antibodies of four adult rabbits [9]. Avian antibodies do not bind to the rheumatoid factor or the mammal's Crystallizable Fragment (Fc) receptor, and, they do not activate mammalian complement system. Altogether, these findings made IgY an attractive possibility to avoid the false positive results to serologic diagnostic.

As a quality immune biologic reagent, IgY antibody presents specificity and sensitivity against immunogens such as native proteins, recombinant proteins, peptides, and haptens. Avian antibodies are found in high titles in the egg yolk because they have there, 2-fold the blood circulating concentration [6]. Furthermore, the translocation of IgY, from blood to egg yolk has a circadian rhythm. In addition, IgY has an equivalent affinity maturity to mammals' IgG antibodies [9]. Regarding the antibodies' antigen binding repertoire, polyclonal and monoclonal IgY antibodies have been produced for a wide range of antigenic targets [11,12].

IgY-Technology applied to domestic animals' diseases

The animals' immune response is featured by cellular and humoral immune response and both are the arms of the acquired immune response [13,14]. In mammals' humoral immune response, the IgM antibodies are predominant in the first exposure to the pathogens whilst the IgG antibodies are commonly produced in subsequent exposures [15]. Regarding the serologic diagnostic platforms, these post-pathogens contact blood circulating antibodies can be named "primary antibodies" and they binding to immobilized antigens attached on the diagnostic platforms forming the immune complexes [16]. The detection these immune complexes is primarily the function of immunodiagnostic approaches. In this context, appropriate tools for an immune complexes' detection have been demonstrated by heterologous, polyclonal or monoclonal antibodies, prepared against the host antibodies [9]. These heterologous antibodies are named secondary antibodies and they are featured by detecting class-specific and, also, specie-specific antibodies [17].

Those secondary antibodies are produced using a different specie host from the target specie. Frequently, the antibodies producing hosts are mammals such as rabbits, goats or mice. Thus, the major class of mammal' antibodies is the IgG, which is the same used as a secondary antibody detecting immune complexes. IgG antibody has been prepared as a high avidity biological reagent which has been conjugated with enzymes, fluorophores or metal nanoparticles, and they are suitable tools amplifying the presence of immune complexes detection [12].

At the 80's decade, the increased attention to animal welfare guided the proposition for an alternative source of IgG secondary antibodies. The IgG production includes the animal's immunization and bleeding for extraction of its blood antibodies. In this context, hens' egg yolk has been introduced as a source of high quality antibodies that have an extraction method free of hens' bleeding procedures [6].

The major egg yolk antibody is the IgY and it has equivalent immune properties to the IgG antibody [9]. IgY is the ancestor of mammal's IgG and it has two antigen binding sites as the IgG and, also, the whole IgY molecule is composed by light and heavy peptide chains. However, the IgG's hinge region is absent in IgY antibodies and they have a lengthy Crystallizable Fragment (Fc) than IgG [15]. In this section, we present the applications of IgY antibodies by IgY-technology approaches diagnosing animals' infectious diseases specially in bovine and equine.

Viral diseases

Bovine and equine herds are susceptible to viral infectious diseases, some of them have a zoonotic potential, and others have a negative impact in the country's agriculture system. For instance, rabies is a worldwide zoonotic disease and affects bovine and equine herds in the rural zone of many countries. Foot and Mouth

Disease Virus (FMDV) is a bovine vesicular disease that causes economy losses. In this context, the Infectious Equine Anemia (IEA) and equine Herpesvirus are financial troubles for equine herds.

Rabies

Rabies is a lethal zoonotic disease that infects mammals as bovines, equines and bats [18,19]. The rabies' control program includes periodically monitoring of wild reservoirs such as bats, massive herd vaccination and bats' population control at risk zones [19]. Rabies infection is confirmed by histological assays as Immune HistoChemistry (IHC) or immune peroxidase antigen detection [20]. In the direct antigen detection assay, cytoplasm Lyssavirus inclusions are detected by specific polyclonal IgG anti-rabies antibodies [21]. IgY polyclonal anti-rabies antibodies has been introduced as a promising immune biological for direct detection of Lyssavirus [22]. In addition, IgY antibodies are properly conjugated with peroxidase and fluorescein [23, 24]. Additionally, after the pepsin digestion, IgY antibodies produce two antigen binding fragments (Fab), however IgG antibodies are degraded into F(ab')₂ [25]. This structural difference, in pepsin-degraded antibodies, has been proposed to influence the tissues' antibody distribution in treatments in animal models [26]. Regarding the IgY use for rabies therapy, it has been suggested that it has higher immunogenicity than the IgG for mammals [27].

Foot and Mouth Disease Virus

Bovine vesicular diseases are very important for livestock because of foot-and-mouth disease caused by Foot and Mouth Disease Virus (FMDV) [28]. FMDV's control is based on vaccination of susceptible animals as bovine herds, for example. The differentiation from vaccinated bovines to FMDV infected bovines is crucial to monitor the herd' health status in a country during a FMD eradication program. Enzyme-linked immune sorbent assay (ELISA) sandwich and a competitive ELISA has been suggested to the FMDV detection [29]. In this context, mammal-based recombinant anti body fragments, like single-chain Fragment Variable (scFv), has been applied in FMDV ELISA platforms [30]. Based on the repertoire of antibody genes, SCFV (Single-Chain Fragment Variable) immune libraries have been constructed from antibody genes of B cells derived from immunized animals [31]. Hen's single-chain antibody fragment libraries are useful to diagnostic and research of biological reagents. The use of avian immunoglobulin genes simplifies the construction of such repertoires since far fewer primer sets are required to access the avian antibody genes of B cell than in mammals [32].

Infectious Equine Anemia Virus

Infectious Equine Anemia Virus (IEAV) is the cause of a disease with morbidity and mortality that infects members of Equidae. In Brazil, officially approved Agar Gel Immuno Diffusion assay (AGID) has been used to detect seropositive animals. Moreover, ELISA test and ImmunoBlot (IB) has been used for epidemiological studies [33]. Phylogenomic analysis suggested that the Brazilian EIAV forms a cluster with strains from the United States [34]. The capsid

protein (p26) is one the major immunogenic proteins during EIAV infection and it is widely used for detection of its specific antibodies. Antigen Capture ELISA (AC-ELISA), based on mammals' monoclonal antibodies, has been reported for quantification of the EIAV's p26 protein level [35]. Polyclonal IgY-based detection antibody has been developed to the human severe acute respiratory syndrome-associated coronavirus (SARS-CoV) nucleocapsid protein for the AC-ELISA diagnosing test in which this avian antibody increased the sensitivity of the system [36].

Bacterial diseases

In this section we shall try to describe perspectives for the IgY antibody technology use in zoonotic bacterial diseases; bovine brucellosis, bovine tuberculosis and equine glanders. Bovine brucellosis is caused by *Brucella abortus* that negatively impacts livestock productivity in Brazil. For this reason, in 2001, Brazil launched a new national program aiming the eradication of animal brucellosis. A recently large-scale study was published and revealed important differences in herd prevalence among Brazilian regions. Regarding brucellosis epidemiology herd size and kind of bovine herd were considered significant risk factors [37]. Bovine tuberculosis is caused by *Mycobacterium bovis* which offers risk of infection to human consumers' of raw milk and homemade fresh cheese in Brazil [38]. Glanders is a highly contagious and often fatal zoonotic disease caused by *Burkholderia mallei* primarily often of solipeds and, already used as a biological weapon [39]. Glanders has regained the status of a re-emerging disease because of the numerous recent out breaks and the pre-symptomatic, or carrier animals, that are potential sources of infection and play a crucial role in the spreading of the infectious agent [40].

Bovine Brucellosis

Bovine brucellosis' diagnosing test has been improved to avoid false positive or false negative results. Recently it was proposed a bovine brucellosis diagnostic bead-based immunoassay by Luminex assay using mammal's monoclonal anti bodies against smooth LipoPolySaccharide (LPS) from *B. abortus* [41]. *B. abortus*' LPS has been the target for immune chromatographic test in bovine brucellosis diagnosing [42]. On a strip, immobilized antibodies bind to bacterial LPS and report an infected animal at field conditions. Other cell wall polysaccharides (M polysaccharide) of *B. abortus* were immunogenic for mammalian antibodies eliminating the cross reactivity in serologic diagnostic therefore, used as a biological reagent for bovine brucellosis' diagnostic [43]. Regarding the bacterial polysaccharide immunogenicity, LPS was a target to produce promising IgY antibodies that were used as a LPS-binding tool for endotoxemia's model therapy [44]. Also, polyclonal IgY against LPS from *B. abortus* have been prepared and conjugated with immune magnetic beads to detect the bacterial antigen [45]. Recently, transplastomic plants were transformed with high immunogenic *B. abortus* protein and then, these antigens from the bacterial-recombinant plants were extracted and used to produce high quality polyclonal IgY against *B. abortus* [46].

Bovine tuberculosis

Intradermal test, bacterial culture and molecular methods such as Polymerase Chain Reaction (PCR) and genome sequencing have been used to diagnose *Mycobacterium bovis* infected animals [47]. However, a sandwich ELISA-based interferon gamma (IFN- γ) detection assay has been suggested as a promising complementary method of diagnosing early stages of *M. bovis* infection [48]. Corroborating this findings, positive samples in the tuberculin skin test are, also, positive in the IFN- γ test [49]. Recently, it was developed a novel immune chromatographic-based test for *M. bovis* detection from fecal samples of reservoir wild animals by polyclonal and monoclonal IgG antibodies against *M. bovis* antigen [50]. In the context of *Mycobacterium* genus, specific and sensitive polyclonal IgY antibodies has been used as an immunobiological reagent for the capture of *Mycobacterium avium* antigen and attesting of the bacterial growth in liquid culture [51]. Also, FITC-labeled and immune magnetic beads-coated polyclonal IgY antibodies against *M. Avium* Paratuberculosis (MAP) enhance the bacterial diagnostic and avoid unspecific reactivity to MAP [51,52]. Engineered IgY antibodies from chicken scFv (“gallibodies”) with a more specific antigen binding fragment have been suggested as a promising tool for sandwich ELISA or immune chromatographic test detecting antigens of *M. bovis* [53].

Equine glanders

PCR-based methods have been used for diagnostic and differentiation among *Burkholderia* genus bacteria [54]. However, recombinant antigen-based immunodiagnostic such as ELISA is highlighting the Glanders’ serologic diagnostic [55]. In addition, indirect ELISA or competitive ELISA prepared with recombinant antigens from *B. mallei* were promising diagnosing equine glanders compared to the Complement Fixation Test (CFT) [56,57]. Commercially CFT has been applied to detection of specific antibodies in seropositive Equidae infected with *B. mallei*, however there are false positive tests and divergences in CFT results from different laboratories [58]. Western Blot test (WB) has been prepared to overcome the false positive results caused by high sensitive CFT, and, hence, WB must be considered a confirmatory test while CFT must be used as a routine test for glanders diagnostic in non-endemic areas [59].

Scientific effort has been focused in production of mammal’s monoclonal antibodies with bactericidal activity against *B. mallei* and, in this context, bacterial LPS was a better target for antibody binding than bacterial proteins, and these antibodies result in a superior protection after experimental infection [60]. A valuable strategy diagnosing *B. mallei* are the scFv antibodies from a phage-displayed libraries and these scFv has been demonstrated to have high specificity to discriminate among the pathogenic *Burkholderia* genus [61]. At the moment, there are not any published research with IgY antibodies and glanders. However, these avian antibodies have demonstrated potential for application in ELISA tests,

monoclonal and scFv antibodies approaches. For this reason, our group is developing IgY anti bodies for application in equine glanders’ diagnostic.

Conclusion

Bovine and equine herds are susceptible to infectious diseases that have zoonotic potential, and hence, they have received attention by the government authorities. Diagnostic approaches have been developed to increase specificity and sensitivity and to detect infected animals. ELISA, immune chromatographic test, fluorescent assays are improving continuously and all of them demand high quality antibodies. Overcoming false positive results, mammals’ antibodies have been prepared as monoclonal anti bodies or scFv from phage-displayed libraries. In this context, avian antibodies such as IgY antibodies are promising high-quality antibodies. The results of IgY research demonstrated the robustness of this immune biological reagent and, also, suggested a wide range for IgY applications in viral and bacterial diseases in bovine and equine. We believe IgY-technology is a new frontier for biological reagents applied to diagnostic approaches in veterinary medicine.

References

- Kaiser P (2012) The long view: a bright past, a brighter future? Forty years of chicken immunology pre- and post-genome. *Avian Pathol* 41(6): 511-518.
- Ibrah ME, Perelman B, Finger A, Uni Z (2017) The role of the bursa of Fabricius in the immune response to vaccinal antigens and the development of immune tolerance in chicks (*Gallus domesticus*) vaccinated at a very young age. *Poult Sci* 96(1): 51-57.
- Boehm T, Hess I, Swann JB (2012) Evolution of lymphoid tissues. *Trends Immunol* 33(6): 315-321.
- Lanning DK, Knight KL (2015) Diversification of the Primary Antibody Repertoire by AID-Mediated Gene Conversion. *Results Probl Cell Differ* 57: 279-293.
- Sepahi A, Salinas I (2016) The evolution of nasal immune systems in vertebrates. *Mol Immunol* 69 : 131-138.
- He JX, Thirumalai D, Schade R, Zhang XY, et al. (2014) Chronobiological studies of chicken IgY: monitoring of infradian, circadian and ultradian rhythms of IgY in blood and yolk of chickens. *Vet Immunol Immunopathol* 160 (3-4): 266-272.
- Meunier M, Guyard-Nicodème M, Vigouroux E, Poezevara T, Beven V, et al. (2017) Promising new vaccine candidates against *Campylobacter* in broilers. *PLoS One* 12(11): e0188472.
- Arnold KB, Chung AW (2018) Prospects from systems serology research. *Immunology* 153 (3) 279-289.
- Ferreira Júnior Á, Santiago FM, Silva MV, Ferreira FB, Macêdo Júnior AG, et al. (2012) Production, characterization and applications for *Toxoplasma gondii*-specific polyclonal chicken egg yolk immunoglobulins. *PLoS One* 7 (7): e40391.
- Leandro NM, Ali R, Koci M, Moraes V, Eusebio-Balcazar PE, et al. (2011) Maternal antibody transfer to broiler progeny varies among strains and is affected by grain source and cage density. *Poult Sci* 90(12): 2730-2739.
- Nie G, Wang T, Lu S, Liu W, Li Y, et al. (2014) Detection of *Clonorchis sinensis* circulating antigen in sera from Chinese patients by immunomagnetic bead ELISA based on IgY. *PLoS One* 9(12): e113208.

12. Da Rocha DG, Fernandez JH, de Almeida CMC, da Silva CL, Magnoli FC, et al. (2017) The complementarity-determining region sequences in IgY antivenom hypervariable regions. *Data Brief* 13: 717-722.
13. Jeurissen SH, Boonstra-Blom AG, Al-Garib SO, Hartog L, Koch G (2000) Defence mechanisms against viral infection in poultry: a review. *Vet Q* 22(4): 204-208.
14. Kaiser P, Howell J, Fife M, Sadeyen JR, Salmon N, et al. (2008) Integrated immunogenomics in the chicken: deciphering the immune response to identify disease resistance genes. *Dev Biol (Basel)* 132: 57-66.
15. Zhang X, Calvert RA, Sutton BJ, Doré KA (2017) IgY: a key isotype in antibody evolution. *Biol Rev Camb Philos Soc* 92(4): 2144-2156.
16. Zhang X, Diraviyam T, Li X, Yao G, Michael A (2016) Preparation of chicken IgY against recombinant E2 protein of bovine viral diarrhoea virus (BVDV) and development of ELISA and ICA for BVDV detection. *Biosci Biotechnol Biochem* 80(12): 2467-2472.
17. Borges IP, Silva MF, Santiago FM, de Faria LS, Júnior ÁF, et al. (2018) Antiparasitic effects induced by polyclonal IgY antibodies anti-phospholipase A2 from *Bothrops pauloensis* venom. *Int J Biol Macromol* 112 : 333-342.
18. Pereira AS, Casseb LMN, Barbosa TFS, Begot AL, Brito RMO, et al. (2017) Rabies Virus in Bats, State of Pará, Brazil, 2005-2011. *Vector Borne Zoonotic Dis* 17(8): 576-581.
19. Singh R, Singh KP, Cherian S, Saminathan M, Kapoor S, et al. (2017) Rabies - epidemiology, pathogenesis, public health concerns and advances in diagnosis and control: a comprehensive review. *Vet Q* 37(1): 212-251.
20. Rahmadane I, Certoma AF, Peck GR, Fitria Y, Payne J, et al. (2017) Development and validation of an immunoperoxidase antigen detection test for improved diagnosis of rabies in Indonesia. *PLoS Negl Trop Dis* 11 (11): e0006079.
21. Bassuino DM, Konradt G, Cruz RA, Silva GS, Gomes DC, et al. (2016) Characterization of spinal cord lesions in cattle and horses with rabies: the importance of correct sampling. *J Vet Diagn Invest* 28(4) : 455-460.
22. Motoi Y, Inoue S, Hatta H, Sato K, Morimoto K, et al. (2005) Detection of rabies-specific antigens by egg yolk antibody (IgY) to the recombinant rabies virus proteins produced in *Escherichia coli*. *Jpn J Infect Dis* 58 (2): 115-118.
23. Nafea NM, Sabbah MA, Al-Suhail R, Mahdavi AH, Asgary S, et al. (2015) Development of hen antihepatitis B antigen IgY-based conjugate for ELISA assay. *Adv Biomed Res* 4: 100.
24. Sert M, Cakir Koc R, Budama Kilinc Y (2017) Novel FITC-Labeled IgY Antibody: Fluorescence Imaging *Toxoplasma Gondii* In Vitro. *Sci Rep* 7(1): 852.
25. Sun S, Mo W, Ji Y, Liu S (2001) Preparation and mass spectrometric study of egg yolk antibody (IgY) against rabies virus. *Rapid Commun Mass Spectrom* 15(9): 708-712.
26. Sevcik C, Salazar V, Díaz P, D'Suze G, Vázquez H (2013) High resolution fluorescence microscopy evidence on the transport of immunoglobulins. Differences between mammalian IgG, F(ab')₂ and avian IgY. *Toxicon* 63: 7-18.
27. Sevcik C, D'Suze G, Salazar V, Díaz P, Vázquez H (2012) Horse IgG- and ostrich IgY-F(ab')₂ groups have different affinities for mice erythrocytes and lymphocytes. Implications for avian immunoglobulin therapeutic usefulness. *Toxicon* 60(6): 1215-1221.
28. Laguardia-Nascimento M, Sales ÉB, Gasparini MR, de Souza NM, da Silva JA, et al. (2016) Detection of multiple viral infections in cattle and buffalo with suspected vesicular disease in Brazil. *J Vet Diagn Invest* 28 (4): 377-381.
29. Sharma GK, Mahajan S, Matura R, Subramaniam S, Mohapatra JK, et al. (2014) Production and characterization of single-chain antibody (scFv) against 3ABC non-structural protein in *Escherichia coli* for sero-diagnosis of Foot and Mouth Disease virus. *Biologicals* 42(6): 339-345.
30. Sridevi NV, Shukra AM, Neelakantam B, Anilkumar J, Madhanmohan M, et al.(2014) Development of anti-bovine IgA single chain variable fragment and its application in diagnosis of foot-and-mouth disease. *Eur J Microbiol Immunol (Bp)* 4(1): 34-44.
31. Ahmad ZA, Yeap SK, Ali AM, Ho WY, Alitheen NB, et al. (2012) scFv antibody: principles and clinical application. *Clin Dev Immunol* 2012: 980250.
32. Chiliza TE, Van Wyngaardt W, Du Plessis DH (2008) Single-chain antibody fragments from a display library derived from chickens immunized with a mixture of parasite and viral antigens. *Hybridoma (Larchmt)* 27 (6): 413-421.
33. Oliveira FG, Cook RF, Naves JHF, Oliveira CHS, Diniz RS, et al. (2017) Equine infectious anemia prevalence in feral donkeys from Northeast Brazil. *Prev Vet Med* 140: 30-37.
34. Tigre DM, Brandão CF, de Paula FL, Chinalia FA, Campos GS, et al. (2017) Characterization of isolates of equine infectious anemia virus in Brazil. *Arch Virol* 162(3): 873-877.
35. Hu Z, Chang H, Chu X, Li S, Wang M, et al. (2016) Identification and characterization of a common B-cell epitope on EIAV capsid proteins. *Appl Microbiol Biotechnol* 100(24): 10531-10542.
36. Palaniyappan A, Das D, Kammila S, Suresh MR, Sunwoo HH (2012) Diagnostics of severe acute respiratory syndrome-associated coronavirus (SARS-CoV) nucleocapsid antigen using chicken immunoglobulin Y. *Poult Sci* 91 (3): 636-642.
37. de Alencar Mota ALA, Ferreira F, Ferreira Neto JS, Dias RA, Amaku M, et al. (2016) Large-scale study of herd-level risk factors for bovine brucellosis in Brazil. *Acta Trop* 164: 226-232.
38. Belchior AP, Lopes LB, Gonçalves VS, Leite RC (2016) Prevalence and risk factors for bovine tuberculosis in Minas Gerais State, Brazil. *Trop Anim Health Prod* 48(2): 373-378.
39. Van Zandt, KE, Greer MT, Gelhaus H C (2013) Glanders: an overview of infection in humans. *Orphanet J Rare Dis* 8: 131.
40. Khan I, Wieler LH, Melzer F, Elschner MC, Muhammad G, et al. (2013) Glanders in animals: a review on epidemiology, clinical presentation, diagnosis and countermeasures. *Transbound Emerg Dis* 60(3): 204-221.
41. Silbereisen A, Tamborrini M, Wittwer M, Schürch N, Pluschke G (2015) Development of a bead-based Luminex assay using lipopolysaccharide specific monoclonal antibodies to detect biological threats from *Brucella* species. *BMC Microbiol* 15: 198.
42. Byzova NA, Zherdev AV, Eskendirova SZ, Baltin KK, Unysheva GB, et al. (2012) [Development of immune chromatographic test system for the rapid detection of lipopolysaccharide antigen and cells of causative agent of bovine brucellosis]. *Prikl Biokhim Mikrobiol* 48(6): 653-661.
43. Ganesh NV, Sadowska JM, Sarkar S, Howells L, Mc Given J, et al. (2014) Molecular recognition of *Brucella* A and M antigens dissected by synthetic oligosaccharide glycoconjugates leads to a disaccharide diagnostic for brucellosis. *J Am Chem Soc* 136(46): 16260-16269.
44. Zhen YH, Fang R, Ding C, Jin LJ, Li XY, et al.(2011) Efficacy of specific IgY for treatment of lipopolysaccharide-induced endotoxemia using a mouse model. *J Appl Microbiol* 111(6): 1524-32.
45. Song D, Qu X, Liu Y, Li L, Yin D, et al. (2017) A Rapid Detection Method of *Brucella* with Quantum Dots and Magnetic Beads Conjugated with Different Polyclonal Antibodies. *Nanoscale Res Lett* 12(1): 179.


46. Alfano EF, Lentz EM, Bellido D, Dus Santos MJ, Goldbaum FA, et al. (2015) Expression of the Multimeric and Highly Immunogenic Brucella spp. Lumazine Synthase Fused to Bovine Rotavirus VP8d as a Scaffold for Antigen Production in Tobacco Chloroplasts. *Front Plant Sci* 6: 1170.
47. Bolaños CAD, Franco MMJ, Souza Filho AF, Ikuta CY, Burbano-Rosero EM, et al. (2018) Nontuberculous mycobacteria in milk from positive cows in the intradermal comparative cervical tuberculin test: implications for human tuberculosis infections. *Rev Inst Med Trop Sao Paulo* 60: e6.
48. Rivalde MÁ, Thomas J, Sevilla I, Serrano M, Ortíz JA, et al. (2017) Development and evaluation of an interferon gamma assay for the diagnosis of tuberculosis in red deer experimentally infected with *Mycobacterium bovis*. *BMC Vet Res* 13(1): 341.
49. Keshavarz R, Mosavari N, Geravand MM, Tadayon K, Pajooi RA, et al. (2016) Interferon- γ assay, a high-sensitivity, specific and appropriate method for detection of bovine tuberculosis in cattle. *Int J Mycobacteriol* 5 Suppl (1): S219.
50. Stewart LD, Tort N, Meakin P, Argudo JM, Nzuma R, et al. (2017) Development of a novel immunochromatographic lateral flow assay specific for *Mycobacterium bovis* cells and its application in combination with immunomagnetic separation to test badger faeces. *BMC Vet Res* 13(1): 131.
51. Shin SJ, Anklam K, Manning EJ, Collins MT (2009) Rapid mycobacterial liquid culture-screening method for *Mycobacterium avium* complex based on secreted antigen-capture enzyme-linked immunosorbent assay. *Clin Vaccine Immunol* 16(5): 613-620.
52. Shin SJ, Lee SS, Manning EJ, Collins MT (2017) Production of and applications for a polyclonal IgY diagnostic reagent specific for *Mycobacterium avium* subsp. *paratuberculosis*. *J Microbiol* 47(5): 600-609.
53. Wemmer S, Mashau C, Fehrsen J, van Wyngaardt W, du Plessis DH (2010) Chicken scFvs and bivalent scFv-C(H) fusions directed against HSP65 of *Mycobacterium bovis*. *Biologicals* 38(3): 407-414.
54. Lowe W, March JK, Bunnell AJ, O'Neill KL, Robison RA et al. (2014) PCR-based Methodologies Used to Detect and Differentiate the *Burkholderia pseudomallei* complex: *B. pseudomallei*, *B. mallei*, and *B. thailandensis*. *Curr Issues Mol Biol* 16: 23-54.
55. Pal V, Kumar S, Malik P, Rai GP (2012) Evaluation of recombinant proteins of *Burkholderia mallei* for serodiagnosis of glanders. *Clin Vaccine Immunol* 19(8): 1193-1198.
56. Sprague LD, Zachariah R, Neubauer H, Wernery R, Joseph M, et al. (2009) Prevalence-dependent use of serological tests for diagnosing glanders in horses. *BMC Vet Res* 5: 32.
57. Singha H, Malik P, Goyal SK, Khurana SK, Mukhopadhyay C, et al. (2014) Optimization and validation of indirect ELISA using truncated TssB protein for the serodiagnosis of glanders amongst equines. *Scientific World Journal* 469407.
58. Khan I, Wieler LH, Melzer F, Gwida M, Santana VL, et al. (2011) Comparative evaluation of three commercially available complement fixation test antigens for the diagnosis of glanders. *Vet Rec* 169(19): 495.
59. Elschner MC, Scholz HC, Melzer F, Saqib M, Marten P, et al. (2011) Use of a Western blot technique for the serodiagnosis of glanders. *BMC Vet Res* 7: 4.
60. Zhang S, Feng SH, Li B, Kim HY, Rodriguez J, et al. (2011) In Vitro and In Vivo studies of monoclonal antibodies with prominent bactericidal activity against *Burkholderia pseudomallei* and *Burkholderia mallei*. *Clin Vaccine Immunol* 18(5): 825-834.
61. Zou N, Newsome T, Li B, Tsai S, Lo SC (2007) Human single-chain Fv antibodies against *Burkholderia mallei* and *Burkholderia pseudomallei*. *Exp Biol Med* (Maywood) 232(4): 550-556.
62. Zhang Q, Chen B, Yang P, Zhang L, Liu Y, et al. (2015) Identification and structural composition of the blood-spleen barrier in chickens. *Vet J* 204(1): 110-116.



Creative Commons Attribution 4.0 International License

For possible submissions Click Here

[Submit Article](#)



Approaches in Poultry, Dairy & Veterinary Sciences

Benefits of Publishing with us

- High-level peer review and editorial services
- Freely accessible online immediately upon publication
- Authors retain the copyright to their work
- Licensing it under a Creative Commons license
- Visibility through different online platforms