



# *Escherichia coli* Genome-Scale Metabolic Models Could Guide Construction of Proof-of-Principle Strains



**Bashir Sajo Mienda\***

Department of Microbiology & Biotechnology, Federal University Dutse, Nigeria

\*Corresponding author: Bashir Sajo Mienda, Department of Microbiology & Biotechnology, Faculty of Science, Federal University Dutse, PMB 7156 Ibrahim Aliyu Bypass, Dutse, Jigawa, Nigeria, Email: b.mienda@fud.edu.ng/bsmienda@gmail.com

Submission: 📅 April 16, 2018; Published: 📅 April 26, 2018

## Opinion

*Escherichia coli* genome-scale metabolic models (GEMs) have been published with ability to predict metabolic engineering capabilities that could be consistent with experimental measurements. However, the GEMs have limited scope, and the models are of two types, metabolism models (M-model), and metabolism and gene expression (ME-model) that could guide the constructions of proof-of-principle strains of particularly *E. coli* bacterium that may find applications in metabolic engineering strategies, synthetic biology [1], and beyond. GEMs have been clearly established to be capable of predicting metabolic engineering capabilities and could sometime lead to biological discoveries for missing reactions and/or missing gene functions [2-4]. In addition, systems metabolic engineering has proof useful with the use of GEMs where time consuming experimental trial and error was shortened by predicting engineering strategies using GEMs. Although sometimes prediction could fail to agree with experimental data, but in that circumstances missing knowledge can be uncovered and gaps in the reconstruction can therefore be bridged leading to novel biological discoveries [3,4].

The GEMs that is designated as M-model does not differentiate between isozymes as such its predictive capability is not as accurate as that of ME model, which integrates metabolism and gene expression data. The construction of the former model (M-model) is relatively much easier, as the full protocol has been previously published [5], while the ME model requires additional expertise, as it integrates both metabolism, and gene expression data [6]. These two models could serve as platforms for construction of proof-of-principle strains. A number of proof-of-principle strains were constructed using *E. coli* GEMs for increasing succinic acid production [7-10] and/or other chemicals such as 1,4 butanediol [11]. These strategies used GEMs that are considered M-models, with limited scope and fairly accurate predictive power. What we hope to see in the future is the of extended version of M-model that

could include gene expression data (ME-model) in predicting proof-of-concept studies that could be much more accurate predictive power that is greater than 80%. In conclusion, the M-models of *E. coli* has been extensively used for the construction of proof-of-concept studies, particularly in increasing succinic acid production using a number of carbon sources, including glucose, and glycerol. Because of its limited scope and fair predictive accuracy, M-models are expected to be extended to ME-models for the construction of future proof-of-principle strains not limited to *E. coli* bacterium alone, but also for the forthcoming GEMs of microbial species with varieties of biotechnological applications in the field of medicine, agriculture, environment, industries and probably beyond.

## References

1. Lee SY, Kim HU (2015) Systems strategies for developing industrial microbial strains. *Nat Biotechnol* 33(10): 1061-1072.
2. Mienda BS (2017) Genome-scale metabolic models as platforms for strain design and biological discovery. *J Biomol Struct Dyn* 35(9): 1863-1873.
3. O'Brien EJ, Monk JM, Palsson BO (2015) Using genome-scale models to predict biological capabilities. *Cell* 161(5): 971-987.
4. Aziz RK, Khaw VL, Monk JM, Brunk E, Lewis R, et al. (2015) Model-driven discovery of synergistic inhibitors against *E. coli* and *S. enterica* serovar Typhimurium targeting a novel synthetic lethal pair, *aldA* and *prpC*. *Front Microbiol* 6: 958.
5. Thiele I, Palsson BO (2010) A protocol for generating a high-quality genome-scale metabolic reconstruction. *Nat Protocols* 5(1): 93-121.
6. O'Brien EJ, Lerman JA, Chang RL, Hyduke DR, Palsson BØ (2013) Genome-scale models of metabolism and gene expression extend and refine growth phenotype prediction. *Mol Syst Biol* 9: 693-693.
7. Mienda BS, MS Shamsir, Illias RM (2016) Model-guided metabolic gene knockout of *gnd* for enhanced succinate production in *Escherichia coli* from glucose and glycerol substrates. *Comput Biol Chem* 61: 130-137.
8. Mienda BS, Shamsir MS, Illias R (2015) Model-aided *atpE* gene knockout strategy in *Escherichia coli* for enhanced succinic acid production from glycerol. *J Biomol Struct Dyn* 34(8): 1705-1716.

9. Mienda BS, Shamsir MS, Illias R (2015) Model-assisted formate dehydrogenase-0 (fdoH) gene knockout for enhanced succinate production in Escherichia coli from glucose and glycerol carbon sources. J Biomol Struct Dyn 34(11): 2305-2316.
10. Lee SJ, Lee DY, Kim TY, Kim BH, Lee J, et al. (2005) Metabolic engineering of Escherichia coli for enhanced production of succinic acid, based on genome comparison and in silico gene knockout simulation. Appl Environ Microbiol 71(12): 7880-7887.
11. Yim H, Haselbeck R, Niu W, Pujol BC, Burgard A, et al. (2011) Metabolic engineering of Escherichia coli for direct production of 1,4-butanediol. Nat Chem Biol 7(7): 445-452.



Creative Commons Attribution 4.0  
International License

For possible submissions Click Here

[Submit Article](#)



### Open Access Biostatistics & Bioinformatics

#### Benefits of Publishing with us

- High-level peer review and editorial services
- Freely accessible online immediately upon publication
- Authors retain the copyright to their work
- Licensing it under a Creative Commons license
- Visibility through different online platforms