

# Importance of the Olfactory Environment in the Laboratory Animal Science


Korsós G<sup>1</sup>, Bertram L<sup>1</sup>, Rüllicke T<sup>2</sup>, Marinho B<sup>3</sup> and Fekete S<sup>1\*</sup>

<sup>1</sup>Department of Veterinary Medicine, Hungary

<sup>2</sup>Department of Veterinary Medicine, Vienna

<sup>3</sup>Exeter University, England

ISSN: 2576-9162



**\*Corresponding author:** Fekete S,  
Department of Veterinary Medicine,  
Hungary

**Submission:**  June 07, 2019

**Published:**  June, 10 2019

Volume 6 - Issue 2

**How to cite this article:** Korsós G, Bertram L, Rüllicke T, Marinho B, Fekete S. Importance of the Olfactory Environment in the Laboratory Animal Science. *Appro Poultry Dairy & Vet Sci* 6(2). APDV.000631.2019. DOI: [10.31031/APDV.2019.06.000631](https://doi.org/10.31031/APDV.2019.06.000631)

**Copyright@** Amr F El Karmoty, This article is distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use and redistribution provided that the original author and source are credited.

## Opinion

The 3<sup>rd</sup> R, the refinement is receiving more and more attention. The optimizing of the auditory and olfactory environment of the laboratory animals is essential, because the environmental smells influence the behaviour and occasionally the epigenetics of mammals. According to the characteristics of the smell substances a chemical map is developed in the olfactory bulb. The vomeronasale organ (VNO) has an important role in the pheromone-transferred communication [1]. The VNO is expressing specific major histocompatibility genes, too. In horses and felidae a specific behaviour form, the Flehmen-reaction facilitates pheromone molecules to get the VNO. The intraspecific communication is attained by means of the pheromones, especially in the social and sexual relations. The kairomones are functioning in the communication between species. The interomones are smell compounds influencing the behaviour or physiology of other species, with or without a positive or negative effect for the releaser. Some interomones may decrease the stress of other species [2], having a significance of animal welfare.

The HEPA-filter of rodents' cages does not isolate smells. In rat experiment the smell substance of the fox faeces (2,4,5-trimethyl-thiol=TMT) caused an expressed freezing reaction. The TMT increased the blood ACTH and corticosteron level in rats. In the behaviour of mice, the concentrated, synthetic TMT had only a repellent effect, nevertheless the TMT in fox faeces proved as a real predator stimulus [3]. The irritative smells like tholuol induced fear or aversive behaviour [4]. The epigenetic factors are regulated independently from the base sequence of the DNA and by means of the germ lines, the environment-induced phenotypic changes can be inherited. Based on the inherited fear against the smells, ethological animal models can be outlined and environmentally safe dog-cat relaxant, as well as rodent repellent can be developed [5].

In the present experiment the effects of neutral and irritative smells and the synthetic trimethyl-thiol (TMT) upon the open-field (OF) behaviour of rats, with the background pathophysiology and histology were studied. Twelve ChR (W1) weaned male rats were investigated in five OF test: in the presence of a perfume, 10% and 100% TMT, citronella oil and methyl-hydroxy-analogue (MHA). Hiding during the exposure of diluted TMT lasted longer, than in case of the concentrated TMT and MHA. If no hiding box present, rats spent more time far from the concentrated TMT- source, than from the MHA or perfume. The passive ethological elements were overwhelming under the influence of concentrated TMT and MHA. The grooming was generally rare. Summerizing, the diluted synthetic TMT partly brings about the effect of fox faeces, while the concentrated TMT is a repellent. The citronella represents a light stressor. Freezing has been detected rarely and the final corticosteron values did not differ from the physiological ones and both necropsy and histopathology revealed no pathological alterations. Taken together, the short-term smell effects are not able to develop chronic stress. Our results are close to the findings of Grau et al. [6] who have found an avoidance of mice from the ethanol (neutral chemical substance) as well as from the fox faeces and concentrated TMT. Further studies required to elucidate the long-term effects of smells and to differentiate between irritative and species-related effects of smells.

## References

1. Leinders-Zufall T, Lane AP, Puche AC, Ma W, Novotny MV, et al. (2000) Ultrasensitive pheromone detection by mammalian vomeronasal neurons. *405*: 792-796.
2. Apfelbach R, Blanchard CD, Blanchard RJ, Hayes RA, McGregor IS (2005) The effect of predator odors in mammalian prey species: A review of field and laboratory studies. *Neurosc Biobehav Rev* 29: 1123-1144.
3. Ferrero DM, Lemon JK, Fluegge D, Pashkovski SL, Korzan WJ, et al. (2011) Detection and avoidance of a carnivore odor by prey. *Proc Natl Acad Sci U S A* 108(27): 11235-11240.
4. Saraiva LR, Kondoh K, Ye X, Yoon K, Hernandez M, Buck LB (2016) Combinatorial effects of odorants on mouse behavior. *Proc Natl Acad Sci* 113: E3300-E3306.
5. Sullivan TS, Crump DR, Sullivan D (1988) Use of predator odors as repellents to reduce feeding damage by herbivores. III. Montane and Meadow Voles (*Microtus montanus* and *Microtus pennsylvanicus*). *J Chem Ecol* 14(1): 363-377.
6. Grau C, Leclercq J, Descout E, Teruel E, Bienboire-Frosini C, et al. (2019) Ethanol and a chemical from fox faeces modulate exploratory behaviour in laboratory mice. *Appl Anim Behav Sci* 213: 117-123.

For possible submissions Click below:

[Submit Article](#)