


Inulin-Type Fructans in Poultry Feed

Fu Youli*

Qufu Normal University, China

ISSN: 2576-9162



*Corresponding author: Fu Youli, Qufu Normal University, Qufu 273165, China

Submission: 📅 June 28, 2019

Published: 📅 July 17, 2019

Volume 6 - Issue 2

How to cite this article: Fu Y. Inulin-Type Fructans in Poultry Feed. *Appro Poultry Dairy & Vet Sci* 6(3). APDV.000637.2019. DOI: [10.31031/APDV.2019.06.000637](https://doi.org/10.31031/APDV.2019.06.000637)

Copyright@ Fu Youli, This article is distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use and redistribution provided that the original author and source are credited.

Introduction

Fructans exist as a wide range of oligo- and polysaccharides in many species of bacteria, fungi, and plants [1]. They are classified into different families on the basis of their glycosidic linkages, consisting of (2→1)-linked β-D-fructofuranosyl units such as inulin, or (2→6)-linked β-D-fructofuranosyl units such as levans, or highly branched structures comprised of both (2→1)- and (2→6)-linked β-D-fructofuranosyl units such as graminans [2,3]. Among which, inulin-type fructans are perhaps by far the most widespread and researched current prebiotics.

The European Union has banned all in-feed use of antibiotics from 2006 and the use of antibiotics in feed is gradually reduced (or intense regulation) in other parts of the world. This perspective has stimulated nutritionists and feed manufacturers to search for new, safer alternatives. The primary alternatives studied include acidification of the feed by organic acids, feeding probiotic organisms and feeding prebiotic compounds.

In the 1980s, the possible potential effects of prebiotics in animal feeds was already recognized. Since then, the interest in the use of prebiotics in animal feeds has resulted in extensive research [4].

Unlike other carbohydrates, the inulin-type fructans are indigestible to mammalian enzymes [5], but may be fermented to produce lactic acid [6]. The non-digestible inulin-type fructans are found widely in many vegetable feed and food ingredients and are perhaps the best studied and documented prebiotics in domesticated animals [7].

At hatching, the gastrointestinal tract of broilers is sterile. Immediately, bacteria originating from the mother, the environment or the diet will colonise in the gastrointestinal tract. In case of mother contacts, a diverse microbial population will enter the gastrointestinal tract. As a result, after the first colonisation, bacterial species coming later in time will have greater difficulty colonising (colonisation resistance) than the initial population. Because of the strict separation of generations in broiler chickens, any bacteria from the environment might colonise (e.g. attach to intestinal binding sites or multiply faster than being removed via chyme passage) the intestinal tract. Those feed components that are resistant to enzymatic degradation, such as inulin-type fructans, serve as a substrate for bacterial activity in the intestinal lumen. The interaction between host nutrition and the intestinal microbiota has been clearly illustrated using germ-free animals. Langhout (1998) clearly showed the importance of controlling the activity of the intestinal microbiota to support gut integrity and to avoid

- (i) bacterial overgrowth,
- (ii) reduced nutrient digestibility and
- (iii) reduced production performance [8].

Feeding inulin-type fructans may be a practical strategy for controlling pathogenic bacteria in chickens. Researchers summarised several experiments in which different types of fructans were fed to broilers alone or in combination with a probiotic to evaluate the effect on colonisation of pathogens (i.e. *Salmonella* spp. and *Campylobacter jejuni*) in caeca and on prechilled poultry carcasses [7,9-13]. Researchers concluded that supplementation of inulin-type fructans in combination with competitive exclusion flora may reduce colonisation by the pathogenic bacteria.

In order to effectively supplement inulin-type fructans in animal feed, additional research is also needed to elucidate the mode of action and the relationship between gut microflora, gut and animal health, and performance. Molecular DNA techniques might be helpful in research to gain further insight into the changes occurring in the composition of the gut microflora and the gene expression in gut tissue and relevant organs.

References

- Andrew JC, Paul B, Ian MS (2002) The structure of starch from seeds and leaves of the fructan-accumulating ryegrass, *Lolium temulentum* L. *J Plant Physiol* 159(3): 221-230.
- Praznik W, Spies T (1993) Fructo-oligosaccharides from *Urginea maritima*. *Carbohydr Res* 243(1): 91-97.
- Hincha DK, Zuther E, Hellwege EM, Heyer AG (2002) Specific effects of fructo- and gluco-oligosaccharides in the preservation of liposomes during drying. *Glycobiol* 12(2): 103-110.
- Verdonk JMAJ, Shim SB, Leeuwen PV, Verstegen MWA (2005) Application of inulin-type fructans in animal feed and pet food. *Br J Nutr* 93(Suppl 1): S125-S138.
- Nilsson U, Oste R, Jagerstad M (1988) Cereal fructans: *in vitro* and *in vivo* studies on availability in rats and humans. *J Nutr* 118(11):1325-1330.
- Muller M, Steller J (1995) Comparative studies of the degradation of grass fructan and inulin by strains of *Lactobacillus paracasei subsp. Paracasei* and *Lactobacillus plantarum*. *J Appl Bacteriol* 78(3): 229-236.
- Flickinger EA, Schreijen EMWC, Patil AR, Hussein HS, Grieshop CM, et al. (2003) Nutrient digestibilities, microbial populations, and protein catabolites as affected by fructan supplementation of dog diets. *J Anim Sci* 81(8): 2008-2018.
- Langhout DJ (1998) The role of the intestinal flora as affected by non-starch polysaccharides in broiler chicks. PhD Thesis, Wageningen Agricultural University.
- Flickinger EA, Fahey GC Jr (2002) Pet food and feed applications of inulin, oligofructose and other oligosaccharides. *Br J Nutr* 87(Suppl 2): S297-S300.
- Bailey JS, Blankenship LC, Cox NA (1991) Effect of fructo oligosaccharide on *Salmonella* colonization of the chicken intestine. *Poult Sci* 70(12): 2433-2438.
- Oyarzabal AA, Conner DE (1996) Application of direct-fed microbial bacteria and fructo oligosaccharides for *Salmonella* control in broilers during feed withdrawal. *Poult Sci* 75(2): 186-190.
- Chambers JR, Spencer JL, Modler HW (1997) The influence of complex carbohydrates on *Salmonella typhimurium* colonization, pH, and density of broiler ceca. *Poult Sci* 76(3): 445-451.
- Fukata T, Sasai K, Miyamoto T, Baba E (1999) Inhibitory effects of competitive exclusion and fructo oligosaccharide, singly and in combination, on *Salmonella* colonization of chicks. *J Food Prot* 62(3): 229-233.

For possible submissions Click below:

[Submit Article](#)