


Preliminary Disease Survey of Domestic Ruminants on Farasan Islands, Saudi Arabia

Soares JF¹ and Wronski T^{2*}

¹Be Wild Aid, Portugal


²School of Biological Sciences, UK

ISSN: 2576-9162



*Corresponding author: Wronski T,
School of Biological Sciences, UK

Submission:  September 14, 2021

Published:  October 06, 2021

Volume 8 - Issue 4

How to cite this article: Soares JF and Wronski T, Preliminary Disease Survey of Domestic Ruminants on Farasan Islands, Saudi Arabia. *Appro Poultry Dairy & Vet Sci* 8(4). APDV. 000693. 2021.
DOI: [10.31031/APDV.2021.08.000693](https://doi.org/10.31031/APDV.2021.08.000693)

Copyright@: Wronski T, This article is distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use and redistribution provided that the original author and source are credited.

Mini Review

The Farasan Islands are situated in the Red Sea in southwestern Saudi Arabia Flamand et al. [1]. The climate is arid with low and highly variable annual rainfall Child et al. [2]. Although there are more than 300 islets, only the two largest Farasan Kebir (381km²) and As Saqid (149km²)-are permanently inhabited by humans Flamand et al. [1] and a prevailing fishing culture has led to the absence of intensified livestock keeping. In recent years, tourism on Farasan Islands has grown significantly, leading to an enhanced demand for goat and sheep products, and therefore an increase in livestock numbers. Counts on Farasan Kebir estimated 1330 individuals, corresponding to a density of 3.5 sheep/goat per km² Wronski et al. [3]. In 1988 the Farasan Islands were gazetted as a protected area for the conservation of the archipelago's outstanding marine diversity and for its unique population of Arabian gazelles Child et al. [2]. Arabian gazelles are threatened with extinction in Saudi Arabia and are classified as "vulnerable" by the IUCN Red List (IUCN/SSC, 2017). The current population of *G. arabica* on the Farasan Islands is estimated to be about 700 individuals Islam et al. [4]. The Farasan population is probably the largest in the world, and thus deserves particular conservation attention Lerp et al. [5]. The emergence of zoonotic disease in humans and their domestic livestock and the interface between domestic animals and wildlife populations has become increasingly important in recent years, especially for species of high conservation value [6-8]. Both *Toxoplasma gondii* and *Brucella spp.* are globally distributed zoonotic pathogens that infect a wide range of warm-blooded animals and cause life-threatening infections in fetuses and immune-compromised adult hosts [9]. Both pathogens are also common causes for abortion in goats, sheep and wild ruminants [10,11]. However, only a few reports address the presence of *Toxoplasma gondii* and *Brucella spp.* in wild and domestic animals in Saudi Arabia [12,13]. Worldwide, the wildlife-livestock interface feed, water and mineral sources are often shared between species contributed to the reemergence of brucellosis and toxoplasmosis in livestock and wildlife [14,15]. Although, *Brucella*. cannot multiply outside of their mammalian hosts it can remain viable within the environment for a short time, confined by high temperatures, dryness, and direct exposure to sunlight. The epidemiological importance of environmental contamination as a source of *Brucella spp.* infection in wildlife species therefore depends largely on local conditions. For Farasan and its particularly harsh environment it is assumed that environmental contamination with *Brucella spp.* lasts only for a few days.

Toxoplasma is thought to be transmitted via oocyst ingestion from an environment contaminated with feces of infected feral cats Miller et al. [16]. In a large feral cat population, fecal deposition was estimated at 107 metric tons/year, resulting in 244 billion oocysts shed by feral cats annually [17-19]. Feral cats are common on Farasan Islands and tolerated by the local community. Since feral cats frequently roam the commonly used foraging habitats of wild and domestic ruminants on Farasan Kebir, it can be assumed that gazelles have a similar risk of infection (and thus similar seroprevalence) compared to domestic livestock. Surveys of *Toxoplasma gondii* infections in domestic animals can provide estimates of environmental contamination and of possible effects on ecologically and behaviorally similar wildlife

species [20,21], for example, found similar seroprevalence in three sympatrically occurring canid species in Indiana, USA.

We thus propose that the risk of toxoplasmosis infection in Farasan gazelles can be indirectly inferred by testing domestic sheep and goats. A major concern of wildlife managers is to prevent the introduction of an infectious disease into a wild population, particularly into sensitive ecosystems such as those found on isolated islands. We thus implemented a preliminary disease survey of domestic livestock on Farasan Islands by testing sera from 225 domestic ruminants (62 sheep and 163 goats) for the presence of antibodies against *Toxoplasma gondii* and *Brucella* species. *Toxoplasma* was tested using a commercially available agglutination test, while *Brucella* spp. were tested using a standard qualitative card test (Brucelloslide-test, bioMérieux) and confirmed by ELISA (COMPELISA-400, AHVLA). *T. gondii* seroprevalence was 33.8% (21/62) in sheep and 54% (88/163) in goats. *Brucella* spp. seroprevalence was 0% for domestic sheep and 0.6% (1/163) for goats. The high *T. gondii* seroprevalence indicated a widespread exposure of domestic ruminants to *T. gondii*, and thus is also of epidemiologic significance for the Arabian gazelle population on the archipelago. Established seroprevalence of *Brucella* spp. was low, as predicted for a hyper-arid environment. The Farasan gazelle population is characterized by a relatively high population density [22,23]. Coupled with its limited geographic distribution (mainly on Farasan Kebir), makes the population particularly vulnerable to stochastic events, such as disease outbreaks [24]. The uncontrolled expansion of feral cats on Farasan Kebir has presumably led to the high prevalence of *T. gondii* in domestic livestock, thus resulting in more oocysts in the environment and raising the risk of toxoplasmosis transmission to humans, domestic livestock, and wildlife. Although cats are typically only infected once before gaining immunity, the huge number of free-ranging cats is enough to maintain a large volume of oocysts on the island [25]. Reducing the number of feral cats by reducing the breeding capacity of its population (e.g., spay and neutering) is an important toxoplasmosis prevention measure, which, together with continuous intensive disease surveillance, may help to protect this gazelle population from disease outbreaks that could compromise the success of ongoing conservation efforts. The use of a vaccine for brucellosis control should be also considered, but no brucellosis vaccine that gives satisfactory results in terms of safety and efficacy in wildlife species, is currently available [26]. Excerpts of this article were published in [27-29].

References

1. Flamand JRB, Thouless CR, Tatwany H, Aamode JF (1988) Status of the gazelles of the Farasan Islands, Saudi Arabia. *Mammalia* 52: 608-610.
2. Child G, Grainger J (1990) A system plan for protected areas for wildlife conservation and sustainable rural development in Saudi Arabia Appendix II. National Commission for Wildlife Conservation and Development.
3. Wronski T (2013) Population development of Arabian Gazelles (*Gazella arabica*) on the Farasan Islands, Saudi Arabia. *Zool Mid East* 59(3): 189-195.
4. Islam MZ, Boug A, Hashim HI, Sher Shah M, Al Subai H, et al. (2018) Adaptation, survival and challenges of released endemic Farasan mountain gazelles on Farasan Islands, Southern Red Sea, Saudi Arabia. In: Soorae (Ed.), *Global Reintroduction Perspectives: Case studies from around the globe*. IUCN/SSC Reintroduction specialist group, gland, switzerland and environment agency, Abu Dhabi, UAE, pp: 144-149.
5. Lerp H, Plath M, Wronski T, Malczyk A, Riesch RR, et al. (2014) Utility of island populations in reintroduction programs-relationships between Arabian gazelles (*Gazella arabica*) from the Farasan Archipelago and endangered mainland populations. *Mol Ecol* 23(8): 1910-1922.
6. Cleaveland S, Hess GR, Dobson AP, Laurenson MK, McCallum, et al. (2002) The role of pathogens in biological conservation. In: Hudson PJ, Rizzoli A, Grenfell BT, Heesterbeek H, Dodson AP (Eds.), *The ecology of wildlife diseases*. Oxford University Press, UK, pp: 139-149.
7. Conrad PA (2007) Detection of *toxoplasma gondii*-like oocysts in cat feces and estimates of the environmental oocyst burden. *J Am Vet Med Assoc* 231(11): 1676-1684.
8. Mazet J, Clifford D, Coppolillo P, Deolalikar A, Erickson J et al. (2009) "One Health" approach to address emerging zoonoses: the HALI Project in Tanzania. *PLoS Med* 6: e1000190.
9. Montoya JG, Liesenfeld O (2004) Toxoplasmosis. *Lancet* 363(9425): 1965-1976.
10. Soares JF, Desta FS, Hanoosh N, Macasero W (2014) Management implications and eradication of *Brucella melitensis* in two different captive gazelle collections from two different gazelle species in the kingdom of Saudi Arabia. *Proceedings of an International Symposium: Disease of Zoo and Wild Animals*, Warsaw, Poland.
11. Tenter AM, Heckeroth AR, Weiss LM (2000) *Toxoplasma gondii*: from animals to humans. *Int J Parasitol* 30(12-13): 1217-1258.
12. Soares JF, Pereira HMA, Desta FS, Macasero W (2013) Mortality of captive sand gazelles (*Gazella gazella*) at king khalid research wildlife center, Saudi Arabia 1988-2011. *Proceedings of an International Symposium: Diseases of Zoo and Wild Animals*, Austria.
13. Mohammed OB, Hussein HB (1994) Antibody prevalence of toxoplasmosis in Arabian gazelles and oryx in Saudi Arabia. *J Wildlife Dis* 30(4): 560-562.
14. Rhyan JC (2000) Brucellosis in terrestrial wildlife and marine mammals. In: Brown C, Bolin C (Eds.), *Emerging diseases of animals*. ASM Press, Washington, USA, pp: 161-184.
15. Van Campen H, Rhyan J (2010) The role of wildlife in diseases of cattle. *Vet Clin N Am Food A* 26(1): 147-161.
16. Miller NL, Frenkel JK, Dubey JP (1972) Oral infections with toxoplasma cysts and oocysts in felines, other mammals, and in birds. *J Parasitol* 58(5): 928-937.
17. Ruiz A, Frenkel JK (1980) *Toxoplasma gondii* in Costa Rican cats. *American J Trop Med Hyg* 29(6): 1150-1160.
18. Dabritz HA, Atwill ER, Gardner IA, Miller MA, Conrad PA (2006) Outdoor fecal deposition by free-roaming cats and attitudes of cat owners and nonowners toward stray pets, wildlife, and water pollution. *J Am Vet Med Assoc* 229(1): 74-81.
19. Dabritz HA, Conrad PA (2010) Cats and toxoplasma: implications for public health. *Zoonoses Public Hlth* 57(1): 34-52.
20. Fredebaugh SL (2010) Habitat overlap and seroprevalence of *Toxoplasma gondii* in wildlife and feral cats in a natural area. University Illinois, USA.
21. Dabritz HA, Miller MA, Atwill ER, Gardner IA, Leutenegger CM, et al. (1999) *Toxoplasma gondii* antibodies in naturally exposed wild coyotes, red foxes, and gray foxes and serologic diagnosis of toxoplasmosis in red foxes fed *T. gondii* oocysts and tissue cysts. *J Parasitol* 85(2): 240-243.
22. Cunningham PL, Wronski T (2011) Twenty years of monitoring of the Farasan gazelle *Gazella gazella* farasani on the Farasan Islands, Saudi Arabia: an overview. *Oryx* 45(1): 50-55.

23. Wronski T, Alageel K, Ismail K, Islam Z (2012) Aerial and ground survey of Farasan mountain gazelles (*Gazella gazella*) on Farasan Islands (June 2012) Quarterly Report July-September 2012, Khalid Wildlife Research Centre, Thumamah, Saudi Arabia.
24. Clifford DL, Mazet J, Dubovi EJ, Garcelon DK, Coonan TJ, et al. (2006) Pathogen exposure in endangered island fox (*Urocyon littoralis*) populations: implications for conservation management. *Biol Conserv* 131(2): 230-243.
25. Conrad P, Mazet J, Clifford D, Scott C, Wilkes M (2009) Evolution of a transdisciplinary "One Medicine-One Health" approach to global health education at the University of California, Davis. *Prev Vet Med* 92(4): 268-274.
26. Davis DS, Elzer PH (2002) Brucella vaccines in wildlife. *Vet Microbiol* 90(1-4): 533-544.
27. Soares JF, Waap H, Macasero W, Desta FS, Khari HK, et al. (2014) Seroprevalence of *Toxoplasma gondii* in free-ranging small mammals from the Kingdom of Saudi Arabia. Proceedings of an International Symposium: European Wildlife Disease Association, Edinburgh, Scotland.
28. Soares JF, Wronski T (2021) Idmi and other mammalian wildlife in the Farasan Islands Protected Area. In: Jawad LAJ (Ed.), *The Arabian Seas: Biodiversity, Environmental Challenges and Conservation Measures*. Springer International Publishing, Switzerland, pp: 1323-1355.
29. IUCN/SSC Antelope Specialist Group (2017) *Gazella arabica*. The IUCN Red List of Threatened Species 2017: e.T117582065A88018124.

For possible submissions Click below:

[Submit Article](#)