

# The Use of Biofertilizers to Increase the Bioactive Compounds in Legumes and Non-Legumes

ISSN: 2637-7802



**\*Corresponding author:** Luís R Silva, CICS-UBI-Health Sciences Research Centre, University of Beira Interior, Covilhã, Portugal

**Submission:**  June 13, 2022

**Published:**  June 17, 2022

Volume 2 - Issue 4

**How to cite this article:** Luís RS\*, Ana CG, José DFF. The Use of Biofertilizers to Increase the Bioactive Compounds in Legumes and Non-Legumes. Biodiversity Online J. 2(4). BOJ. 000545. 2022.

**Copyright@** Luís RS. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use and redistribution provided that the original author and source are credited.

**Luís RS<sup>1,2\*</sup>, Ana CG<sup>1</sup> and José DFF<sup>1</sup>**

<sup>1</sup>CICS-UBI-Health Sciences Research Centre, University of Beira Interior, Portugal

<sup>2</sup>Center of Potential and Innovation of Natural Resources, Polytechnic Institute of Guarda, Portugal

## Opinion

Increasing interest in preserving the environment and consumer health is leading to changes in production methods and dietary habits. Consumers are increasingly demanding safe functional foods with health-promoting properties focused primarily on protection against carcinogenesis and oxidative processes. The consumption of fresh fruits and vegetables containing bioactive compounds has increased significantly in recent years and numerous studies have been conducted on the potential benefits of such compounds for various aspects of human health [1]. The increase in quality demanded by consumers requires specific agricultural practices, such as fertilization with products containing microorganisms [2]. Inoculation with Plant Growth-Promoting Rhizobacteria (PGPR), which reduce the chemicals added to plants, can improve plant growth through various direct mechanisms such as nitrogen fixation, phosphate solubilization, and the production of various compounds such as phytohormones or indirect mechanisms such as the production of siderophores [3]. In addition, the quality of plants and fruits can be improved these microorganisms can mobilize nutrients for plants, influence the hormonal balance of plants or produce microbial inhibitors that prevent the growth of phytopathogens [1,3-5].

To achieve both goals of promoting plant growth in legumes and non-legumes and human health benefits, it is necessary to use non-pathogenic microorganisms in biofertilization programs [6]. Human nutrition is a serious problem nowadays due to various factors including economic, energy and political factors. In developing countries, legumes and cereals are the main, and sometimes the only one sources of proteins for human nutrition [7]. Legumes have the ability to fix atmospheric nitrogen by establishing symbioses with nitrogen-fixing bacteria, called rhizobia, which can convert atmospheric di-nitrogen into ammonia that is available for plants [4,7-9]. Inoculation of a legume with rhizobia leads to metabolic changes in the plant, of which the most studied to date are the increase in nitrogen and protein content and has been used in agriculture to improve the yield of various legumes, reducing the use of chemical allowing health and environmental protection [10]. In recent, the study of other legume components more related to the human health has gained interest. In addition to high-quality proteins, legumes contain several bioactive compounds such as polysaccharides, bioactive peptides, phenolics, including isoflavones, carotenoids, tocopherols and fatty acids, as well as other phytochemicals, which make legumes excellent functional foods and include them in nutraceutical products [11]. The studies on the effect of rhizobia inoculation on legume

bioactive compounds have been conducted in edible legumes, with particular emphasis on the studies conducted in soybean, given the economic importance of this legume worldwide. Couto [9] reported that inoculation of *Glycine max* with *Bradyrhizobium japonicum* increased the content of phenolic compounds and organic acids. The same was observed for volatile compounds, although compounds such as linalyl acetate, menthyl acetate and  $\alpha$ -farnesene were detected only in inoculated plants. The phenolic extracts showed stronger antioxidant capacity than the extracts from organic acid. In addition, extracts from plants inoculated with *B. japonicum* exhibited significantly higher antioxidant activity than control samples. In addition, inoculated seeds had higher total fatty acid content, which was due to an increase in monounsaturated (MUFA) and Polyunsaturated Fatty Acid (PUFA) content [8]. Similar results were observed after inoculation of chickpea (*Cicer arietinum*) with a *Mesorhizobium* strain which did not cause a significant increase in antioxidant potential, but significantly increased the content of flavonoids in the seeds [12]. Considering the economic importance of soybean around the world, experiments on soybean have been highlighted.

Strains of different *Rhizobium* species have also been reported to act as root colonizers and plant growth-promoters in some vegetables, such as pepper (*Capsicum annuum*) [5,13], tomato (*Solanum lycopersicum*), lettuce (*Lactuca sativa*) and carrot (*Daucus carota*) [6,14]. Flores Félix [1] showed that *Phyllobacterium*, phylogenetically related to the *Rhizobiaceae* family, promotes the growth of strawberry, one of the most important berry fruits consumed in Europe. Also, this strain was able to improve the content on vitamin C on fruit, been this compound the main organic acid involved in antioxidant activity of this fruit. In another study with strawberries, Flores Félix [15] investigated the effects of inoculation of strawberry plants with strain PEPV16 of *Rhizobium* sp. from the phylogenetic group of *R. legumin* Sarum. The results showed that PEPV16 promoted strawberry growth by significantly increasing the number of stolon's, flowers, and fruits, had higher concentrations of Fe, Zn, Mn, and Mo, and also had higher concentrations of organic acids, such as citric and malic acids, and lower amounts of ascorbic acid than fruits, compared with uninoculated controls. Biofertilization with rhizobacteria can increase the yield and quality of legumes and non-legumes, helping to protect human health and the environment.

## References

- Flores FJD, Silva LR, Rivera LP, Marcos GM, García F, et al. (2015) Plants probiotics as a tool to produce highly functional fruits: the case of *Phyllobacterium* and vitamin C in strawberries. *PLoS One* 10(4): e0122281.
- Pascual I, Azcona I, Aguirreola J, Morales F, Corpas FJ, et al. (2010) Growth, yield, and fruit quality of pepper plants amended with two sanitized sewage sludges. *J Agric Food Chem* 58(11): 6951-6959.
- Lugtenberg B, Kamilova F (2009) Plant-growth-promoting rhizobacteria. *Annual Review of Microbiology* 63: 541-556.
- Silva LR, Azevedo J, Pereira MJ, Carro L, Velazquez E (2014) Inoculation of the nonlegume *Capsicum annuum* (L.) with *Rhizobium* strains. 1. Effect on bioactive compounds, antioxidant activity, and fruit ripeness. *J Agric Food Chem* 62(3): 557-564.
- Silva LR, Azevedo J, Pereira MJ, Carro L, Velazquez E, et al. (2014) Inoculation of the nonlegume *Capsicum annuum* L. with *Rhizobium* strains. 2. Changes in sterols, triterpenes, fatty acids, and volatile compounds. *J Agric Food Chem* 62(3): 565-573.
- García FP, Carro L, Robledo M, Ramírez BMH, Flores FJD, et al. (2012) *Rhizobium* promotes non-legumes growth and quality in several production steps: towards a biofertilization of edible raw vegetables healthy for humans. *PLoS One* 7(5): e38122.
- Velazquez E, Silva LR, Peix A (2010) Legumes: A healthy and ecological source of flavonoids. *Current Nutrition & Food Science* 6(2): 109-144.
- Silva LR, Pereira MJ, Azevedo J, Mulas R, Velazquez E, et al. (2013) Inoculation with *Bradyrhizobium japonicum* enhances the organic and fatty acids content of soybean (*Glycine max* (L.) Merrill) seeds. *Food Chemistry* 141(4): 3636-3648.
- Couto C, Silva LR, Valentão P, Velázquez E, Peix, A, et al. (2011) Effects induced by the nodulation with *Bradyrhizobium japonicum* on *Glycine max* (soybean) metabolism and antioxidant potential. *Food Chemistry* 127: 1487-1495.
- Silva LR, Bento C, Gonçalves AC, Flores FJD, Ramírez BMH, et al. (2017) Legume bioactive compounds: Influence of rhizobial inoculation. *AIMS microbiol* 3(2): 267-278.
- Silva LR, Alvaro P, Albuquerque C, Velazquez E (2016) Bioactive compounds of legumes as health promoters. *Bentham Science Publisher* 1: 3-27.
- Singh A, Jain A, Sarma BK, Upadhyay RS, Singh HB (2014) Beneficial compatible microbes enhance antioxidants in chickpea edible parts through synergistic interactions. *LWT-Food Science and Technology* 56(2): 390-397.
- Silva LR, Azevedo J, Pereira MJ, Carro L, Velazquez E, et al. (2014) Inoculation of the nonlegume *Capsicum annuum* (L.) with *Rhizobium* strains. 1. effect on bioactive compounds, antioxidant activity, and fruit ripeness. *J Agric Food Chem* 62(3): 557-564.
- Flores FJD, Menéndez E, Rivera LP, Marcos GM, Martínez HP, et al. (2013) Use of *Rhizobium leguminosarum* as a potential biofertilizer for *Lactuca sativa* and *Daucus carota* crops. *Journal of Plant Nutrition and Soil Science* 176(3): 876-882.
- Flores FJD, Marcos GM, Silva LR, Menéndez E, Martínez ME, et al. (2015) *Rhizobium* as plant probiotic for strawberry production under microcosm conditions. *Symbiosis* 67: 25-32.