


Hairy Polyp

George R Wettach*

Washington University, USA

Clinical Image



*Corresponding author: Washington University, USA

Submission: 📅 October 05, 2020

Published: 📅 October 22, 2020

Volume 4 - Issue 3

How to cite this article: George R Wettach. Hairy Polyp. Cohesive J Microbiol Infect Dis. 4(3). CJMI. 000587. 2020. DOI: [10.31031/CJMI.2020.04.000587](https://doi.org/10.31031/CJMI.2020.04.000587)

Copyright@ George R Wettach. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use and redistribution provided that the original author and source are credited.



Figure 1: Hairy Polyp

(Figure 1) A female infant was delivered at 39 weeks gestation by emergent cesarean section to a 27-year-old G3P1. The patient was initially cyanotic and tachypneic. She was admitted to the NICU for ongoing supplemental oxygen. After gagging during attempted feeding, a relatively large (2.7 x 1.9 x 0.7 cm) lesion was retrieved from the oral cavity (Panel A). It had a very narrow stalk and was covered with numerous fine hairs. Histologically, much of the specimen was composed of fibroadipose tissue with a few hair follicles along the periphery. Hairy polyps are rare naso-oropharyngeal tumors usually associated with respiratory and feeding difficulties shortly after birth. Their etiology and precise classification remains uncertain..

For possible submissions Click below:

[Submit Article](#)