


White Matter, Psychomotor Development, Cognitive and Behavior Disorders in Patient with Congenital Heart Defects

Emmanuel Suluba^{1,2*}, Liu Shuwei^{1,2}, Ally Mwanga³ and Balowa Musa³

¹School Basic Medical Sciences, China

²Department of Anatomy and Embryology, China

³School of Medicine, China



***Corresponding author:** Emmanuel Suluba, School Basic Medical Sciences, Shandong, China

Submission: 📅 October 04, 2019

Published: 📅 October 15, 2019

Volume 2 - Issue 4

How to cite this article: Emmanuel Suluba, Liu Shuwei, Ally Mwanga, Balowa Musa. White Matter, Psychomotor Development, Cognitive and Behavior Disorders in Patient with Congenital Heart Defects. COJ Technical & Scientific Research.2(4). COJTS.000541.2019.

Copyright@ Emmanuel Suluba, This article is distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use and redistribution provided that the original author and source are credited.

Abstract

A hemodynamic disturbance in children with congenital heart defects (CHD) impairs cerebral perfusion resulting into brain neural injury. Neural injury in CHDs patients can occur prenatally, postnatal or during surgical repair of the defects (reperfusion injury) often following hypoxia induced oxidative stress. The brain subventricular zone (SVZ) contain postnatal niche of neural stem cells. SVZ cells are very sensitive to hypoxia and may be damaged in children with CHDs. Oligodendriocytes and myelinated fibers are documented to be most vulnerable in conditions of diminished oxygen supply to the brain thus frequently damaged in patient with CHDs. The evaluation of brain associated consequences is frequently overlooked in many cases of CHDs. Therefore, this mini review put forward the impact of CHDs on brain white matter, cognitive and behavior. In addition, it highlights the important infant cognitive evaluation tools and novel techniques used to study the brain white matter.

Keywords: Congenital heart diseases CHDs; White matter injury; Brain volume; Cortical thickness; Psychomotor development; Cognitive; Behavior

Abbreviations: CHD: Congenital Heart Defects; SVS: Sub Ventricular Zone; HLHS: Hypoplastic Left Heart Syndrome; TOF: Tetralogy of Fallot; TGV: Transposition of Greater Vessels

Introduction

The brain is the part of central nervous system which is responsible for coordination and control of all body activities. It consists of grey matter, white matter and neuronal supporting cells. The white matter contains neuronal conducting fibers which transmit information and connect different part of grey matter in the brain. These fibers are classified as; projection fibers, communicating fibers and association fibers [1]. The brain is very sensitive to hypoxia, thus hypoxemic spells can result into permanent neural injury. The damage of brain white matter may affect children psychomotor development, with long term consequences in their cognitive and behavior [2]. The intrauterine and extra uterine hemodynamic disturbance resulting into decrease in cerebral perfusion in fetuses and neonates with CHDs and subsequently affects brain development. Neonates with congenital heart defects have reduced brain oxygen delivery to the brain [3], cause brain maturation delays [4,5] and reduce white matter and grey matter volume [6].

White matter changes cognitive and behavior

The brain subventricular zone (SVZ) contains postnatal niche of neuronal stem cells. Proliferation and neurogenesis of the cells in (SVZ) is impaired by hypoxia [7]. Impairment of regional and global cerebral perfusion therefore affect both structural and function of the brain [8-12] and has detrimental effect on central nervous system [13-17] consequently increase risk of both neurodevelopmental and psychiatric disorders including cognitive, speech, motor and behavioral [18-20]. Children with CHDs have increased risk of hyperactivity and attention deficit [21]. The prenatal brain growth is also impaired in CHDs [22] particularly with CHDs phenotypes namely; hypo plastic left heart syndrome (HLHS), Tetralogy of Fallot (TOF) and transposition of greater vessels (TGV) [23].

White matter injury is particularly common in patient with various forms of CHDs [24-26] with oligodendrocytes and myelinated neurons most affected [27], this strongly correlate with reduced cognitive development [28]. Patient with CHDs have low volume of corpus callosum [29], diffuse microstructure abnormalities in splenium, a part which is involved in visual spatial function [30], uncinate fasciculus and right middle cerebellar peduncle abnormalities which affect memory and auditory attention [31] as well as decreasing thalamocortical development [32]. Diffusion tensor imaging has become the gold standard for diagnosis of brain neurological disorders apart from neurite orientation dispersion and density imaging model which are regarded as more advanced models [33].

DTI provide microstructure assessment of white matter injury within the brain. DTI use the rate of diffusion of water in a tissue to produce images. Using DTI, we can quantitatively measure Fractional Anisotropy, mean diffusivity (MD), axial diffusivity (AD), and radial diffusivity (RD) [34]. Standard infant development evaluation tools such as; Bayley scale of infant development (BSID-II), Denver development screening test (DDST), Mental developmental index (MDI) and psychomotor developmental index (PDI) [35-39] are used to study the impact of white matter injury on psychomotor, cognitive and behavior development in patient with CHDs.

Conclusion

Congenital heart diseases affect both the heart and brain. Indeed, disturbance in fetal hemodynamics which do occur either intra or extra-uterine in CHDs patients have severe brain consequences. We advocate the routine evaluation of white matter using contemporary imaging techniques in order to detect the extent of white matter injury. In addition; we should also pay attention in assessing the psychomotor, cognitive and behavior in CHDs using standard infant development evaluation tools. Therefore, we should not only take care of the heart in management of CHD patients but also the brain. This will ensure proper management for patient with CHDs..

References

- Schmahmann JD, Smith EE, Eichler FS, Filley CM (2008) Cerebral white matter: Neuroanatomy, clinical neurology and neurobehavioral correlates. *Ann N Y Acad Sci* 1142: 266-309.
- Muetzel RL, Mous SE, Vander Ende J, Blanken LM, Vander Lugt A, et al. (2015) White matter integrity and cognitive performance in school-age children: A population-based neuroimaging study. *Neuroimage* 119: 119-128.
- Lim JM, Kingdom T, Saini B, Chau V, Post M, et al. (2016) Cerebral oxygen delivery is reduced in newborns with congenital heart disease. *J Thorac Cardiovasc Surg* 152(4): 1095-1103.
- Licht DJ, Shera DM, Clancy RR, Wernovsky G, Montenegro LM, et al. (2009) Brain maturation is delayed in infants with complex congenital heart defects. *J Thorac Cardiovasc Surg* 137(3): 529-536.
- Shedeed SA, Elfaytouri E (2011) Brain maturity and brain injury in newborns with cyanotic congenital heart disease. *Pediatr Cardiol* 32(1): 47-54.
- Schellen C, Ernst S, Gruber GM, Mlczoch E, Weber M, et al. (2015) Fetal MRI detects early alterations of brain development in tetralogy of fallot. *Am J Obstet Gynecol* 213(3): 392.
- Morton PD, Korotcova L, Lewis BK, Bhuvanendran S, Ramachandra SD, et al. (2017) Abnormal neurogenesis and cortical growth in congenital heart disease. *Sci Transl Med* 9(374).
- Claessens NHP, Kelly CJ, Counsell SJ, Benders MJNL (2017) Neuroimaging, cardiovascular physiology and functional outcomes in infants with congenital heart disease. *Dev Med Child Neurol* 59(9): 894-902.
- Peyvandi S, Donofrio MT (2018) Circulatory changes and cerebral blood flow and oxygenation during transition in newborns with congenital heart disease. *Semin Pediatr Neurol* 28: 38-47.
- Nagaraj UD, Evangelou IE, Donofrio MT, Vezina LG, McCarter R, et al. (2015) Impaired global and regional cerebral perfusion in newborns with complex congenital heart disease. *J Pediatr* 167(5): 1018-1024.
- Watanabe K, Matsui M, Matsuzawa J, Tanaka C, Noguchi K, et al. (2009) Impaired neuroanatomic development in infants with congenital heart disease. *J Thorac Cardiovasc Surg* 137(1): 146-153.
- Cordina R, Grieve S, Barnett M, Lagopoulos J, Malitz N, et al. (2014) Brain volumetric, regional cortical thickness and radiographic findings in adults with cyanotic congenital heart disease. *Neuroimage Clin* 4: 319-325.
- Esquivel HFJ, Mendieta AGG, Pliego RFB, Otero OGA (2015) Electroencephalographic and neurodevelopmental disorders in severe congenital heart disease: A follow-up study. *Gac Med Mex* 151(5): 588-598.
- Yilmaz IZ, Erdur B, Ozbek E, Mese T, Karaarslan U, et al. (2018) Neurodevelopmental evaluation of children with cyanotic congenital heart disease. *Minerva Pediatr* 70(4): 365-370.
- Wernovsky G, Shillingford AJ, Gaynor JW (2005) Central nervous system outcomes in children with complex congenital heart disease. *Curr Opin Cardiol* 20(2): 94-99.
- Donofrio MT, Duplessis AJ, Limperopoulos C (2011) Impact of congenital heart disease on fetal brain development and injury. *Curr Opin Pediatr* 23(5): 502-511.
- Clouchoux C, Du Plessis AJ, Bouyssi KM, Tworetzky W, Mc Elhinney DB, et al. (2013) Delayed cortical development in fetuses with complex congenital heart disease. *Cereb Cortex* 23(12): 2932-2943.
- Nattel SN, Adrianzen L, Kessler EC, Andelfinger G, Dehaes M, et al. (2017) Congenital heart disease and neurodevelopment: Clinical manifestations, genetics, mechanisms and implications. *Can J Cardiol* 33(12): 1543-1555.
- Herberg U, Gürich HH (2012) Neurological and psychomotor development of fetuses and children with congenital heart disease-causes and prevalence of disorders and long-term prognosis. *Z Geburtshilfe Neonatol* 216(3): 132-140.
- Martinez-Biarge M, Jowett VC, Cowan FM, Wusthoff CJ (2013) Neurodevelopmental outcome in children with congenital heart disease. *Semin Fetal Neonatal Med* 18(5): 279-285.
- Shillingford AJ, Glanzman MM, Ittenbach RF, Clancy RR, Gaynor JW, et al. (2008) Inattention, hyperactivity, and school performance in a population of school-age children with complex congenital heart disease. *Pediatrics* 121(4): e759-e767.
- Sun L, Macgowan CK, Sled JG, Yoo SJ, Manlhiot C (2015) Reduced fetal cerebral oxygen consumption is associated with smaller brain size in fetuses with congenital heart disease. *Circulation* 131(15): 1313-1323.
- Hansen T, Henriksen TB, Bach CC, Matthiesen NB (2017) Congenital heart defects and measures of prenatal brain growth: A systematic review. *Pediatr Neurol* 72: 7-18.
- Mulkey SB, Ou X, Ramakrishnaiah RH, Glasier CM, Swearingen CJ (2014) White matter injury in newborns with congenital heart disease: A diffusion tensor imaging study. *Pediatr Neurol* 51(3): 377-383.
- Dimitropoulos A, McQuillen PS, Sethi V, Moosa A, Chau V (2013) Brain injury and development in newborns with critical congenital heart disease. *Neurology* 81(3): 241-248.

26. Miller SP, McQuillen PS, Hamrick S, Xu D, Glidden DV (2007) Abnormal brain development in newborns with congenital heart disease. *N Engl J Med* 357(19): 1928-1938.
27. Morton PD, Ishibashi N, Jonas RA, Gallo V (2015) Congenital cardiac anomalies and white matter injury. *Trends Neurosci* 38(6): 353-363.
28. Rollins CK, Watson CG, Asaro LA, Wypij D, Vajapeyam S (2014) White matter microstructure and cognition in adolescents with congenital heart disease. *J Pediatr* 165(5): 936-944.
29. Hagmann C, Singer J, Latal B, Knirsch W, Makki M (2016) Regional microstructural and volumetric magnetic resonance imaging (MRI) abnormalities in the corpus callosum of neonates with congenital heart defect undergoing cardiac surgery. *J Child Neurol* 31(3): 300-308.
30. Paquette LB, Wisnowski JL, Ceschin R, Pruetz JD, Detterich JA, et al. (2013) Abnormal cerebral microstructure in premature neonates with congenital heart disease. *AJNR Am J Neuroradiol* 34(10): 2026-2033.
31. Brewster RC, King TZ, Burns TG, Drossner DM, Mahle WT (2015) White matter integrity dissociates verbal memory and auditory attention span in emerging adults with congenital heart disease. *J Int Neuropsychol Soc* 21(1): 22-33.
32. Paquette LB, Smith VJK, Ceschin R, Nagasunder AC, Jackson HA, et al. (2015) Abnormal development of thalamic microstructure in premature neonates with congenital heart disease. *Pediatr Cardiol* 36(5): 960-969.
33. Karmacharya S, Gagoski B, Ning L, Vyas R, Cheng HH, et al. (2018) Advanced diffusion imaging for assessing normal white matter development in neonates and characterizing aberrant development in congenital heart disease. *Neuroimage Clin* 19: 360-373.
34. Ranzenberger LR, Snyder T (2019) Diffusion tensor imaging. In *stat pearls*, Stat Pearls Publishing, Treasure Island (FL), Island.
35. Sharp M, De Mauro SB (2017) Counterbalanced comparison of the BSID-II and Bayley-III at eighteen to twenty-two months corrected age. *J Dev Behav Pediatr* 38(5): 322-329.
36. O Shea TM, Joseph RM, Allred EN, Taylor HG, Leviton A, et al. (2018) Accuracy of the Bayley-II mental development index at 2 years as a predictor of cognitive impairment at school age among children born extremely preterm. *J Perinatol* 38(7): 908-916.
37. Borowitz KC, Glascoe FP (1986) Sensitivity of the denver developmental screening test in speech and language screening. *Pediatrics* 78(6): 1075-1088.
38. Lowe JR, Erickson SJ, Schrader R, Duncan AF (2012) Comparison of the bayley II mental developmental index and the bayley III cognitive scale: Are we measuring the same thing? *Acta Paediatr* 101(2): e55-e58.
39. Johnson S, Moore T, Marlow N (2014) Using the bayley-III to assess neurodevelopmental delay: Which cut-off should be used? *Pediatr Res* 75(5): 670-674.

For possible submissions Click below:

[Submit Article](#)