

Possibilities of Speciation Following Anthropogenous Environmental Changes in the Central Sandy Area of the Carpathian Basin through the Example of *Festuca Taxa*

Károly Penksza^{1*}, László Kovács², Zsuzsa Lisztes Szabó³ and Gergely Pápay¹

^{1,2}Szent István University, Faculty of Agricultural and Environmental Sciences, Hungary

³Isotope Climatology and Environmental Research Centre, Institute for Nuclear Research, Hungary

ISSN: 2578-0336



***Corresponding author:** Károly Penksza, Szent István University, Faculty of Agricultural and Environmental Sciences, Institute of Crop Production Sciences, Páter K. u. 1. H-2100 Gödöllő, Hungary

Submission: 📅 October 13, 2020

Published: 📅 March 26, 2021

Volume 8 - Issue 2

How to cite this article: Károly Penksza, László Kovács, Zsuzsa Lisztes Szabó, Gergely Pápay. Possibilities of Speciation Following Anthropogenous Environmental Changes in the Central Sandy Area of the Carpathian Basin through the Example of *Festuca Taxa*. Environ Anal Eco stud. 8(2). EAES. 000683. 2021.

DOI: [10.31031/EAES.2021.08.000683](https://doi.org/10.31031/EAES.2021.08.000683)

Copyright@ Károly Penksza, This article is distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use and redistribution provided that the original author and source are credited.

Abstract

Our goal is to check and revise the dominant *Festuca* species of vegetation types formed under extraordinary conditions through morpho-taxonomic and ploidy analyses. To accomplish this, we had to add examinations of dominant species in grasslands further from the Danube in the Carpathian Basin and East Central Europe. Individuals of examined taxa were analysed using 26 parameters of the inflorescences. Ploidy was analysed using low cytometry. After deforestation and shrubcutting, bare soil patches of areas exposed to anthropogenous effects had provided an opportunity new vegetation to form. As a result of this work, new species *Festuca pseudovaginata* had been discovered here, which is endemic in the Carpathian Basin. Survey continues in order to clear other hardly identifiable taxa. The results have confirmed the presence of the species, but we also have new occurrences discovered.

We verified *F. vaginata* and *F. pseudovaginata* from open sandy areas. In closing grasslands *F. javorkae* and *F. wagneri* appears. In Slovakia we found *F. wagneri* and *F. pseudovaginata* as new species in the country's flora. We could add new appearance data of *F. javorkae* and describe *F. brevipila* as a new taxon of the Hungarian flora. Furthermore, a possibly new species also appeared during our research, on which we found distinctive morphological features, but to describe it as new species it needs further ITS analyses.

Keywords: *Festuca vaginata*; *Festuca psammophila* series; Sandy vegetation

Introduction

Those taxa of *Festuca species* in the Carpathian Basin which have bowed base and stem leaves are identified and mentioned by various authors as belonging to the aggregation of *Festuca ovina* [1-9]. These taxa can be identified easily based on their characteristic tissue structure and molecular genetic analyses [10-16]. Pawlus [17] separated a new series within the *Festuca genus*. This is the *psammophila* series, which includes *F. polesica Zapal*, *F. vaginata W. K.*, *F. psammophila Host.*, *F. pallens*. Subsequently, Šmarda et al. [17] added *F. pseudovaginata Penksza and F. glaucina Stohr.* to it.

The species which were determinate in the original natural vegetation [18] were expected to appear on the new bare soil patches. So, we wanted to answer the question: on new surfaces formed by human interventions, is there a possibility to form new vegetation units and species, or will the natural vegetation and its main species appear?

Material and Methods

The *Festuca* individuals collected in the Carpathian Basin were put through morphological and ploidy examination. Meanwhile, we made coenological records in every type of sandy grasslands along the Danube from Austria to Romania. Furthermore, we set up a collection of living plants, which ensures individuals growing in the same conditions to molecular surveys. We also modelled the appearance, composition and dominant species of new vegetation on bare soil surfaces. The area has been being managed since 2006. The study area lies on the Pest plain near the left bank of the Danube.

Result

Based on our results, we confirmed the appearance of *Festuca vaginata*, *F. pseudovaginata*, *F. javorkae*, *F. wagneri* in the natural grasslands, and discovered new occurrences of *F. brevibila* and *F. javorkae* in the area. In Slovakia, *F. wagneri* appeared as a new species of the flora of the country. *Festuca pseudovaginata* habits only the Pannon region, we found endemic and natural stands of it, but on secondary habitats it was confirmed as a completely new species. Furthermore, taxa of disturbed vegetations are being examined. These habitats are potential hotspots of speciation.

On bare soil surfaces of areas exposed to anthropogenous effects, two species of the genus *Festuca* have become dominant. One of them, *Festuca pseudovaginata* was described as a new species [18] and is endemic in the Carpathian Basin [19]. Examinations are being continued in order to clear hardly identifiable taxa. Our survey has confirmed the occurrence of the species so far, and we discovered new occurrences too.

Acknowledgement

The work was supported by OTKA K-125423. Author LSZ was supported by the European Union and the State of Hungary, co-financed by the European Regional Development Fund in project GINOP-2.3.2.-15-2016-00009 'ICER.'

References

- Adler W, Oswald K, Fischer R (1994) Exkursionsflora von Österreich. Ulmer, Austria.
- Horánszky A, Jankó B, Vida G (1971) Zur Biosystematik der *Festuca ovina*-gruppe in Ungarn. Ann Univ Sci Budapest Sect Biol 13: 95-101.
- Májovszký J (1962) Adnotationes ad species gen. *Festuca* florum Slovakiae additamentum I. Acta Fac Rer Nat Univ Comen 7: 317-355.
- Patzke E (1968) Zur Kenntnis der Sammelart *Festuca ovina* L im südlichen Niedersachsen. Götting Flor Rundbr 4: 14-17.
- Soó R (1973) Zeitgemässe Taxonomie der *Festuca ovina*-gruppe. Acta Bot Sci Hung 18: 363-377.
- Stace CA, Al-Bermani AK, Wilkinson M J (1992) The distinction between the *Festuca ovina* L and *Festuca rubra* L aggregates in the British Isles 19: 7-14.
- Sávulescu T (1972) Flora Republicii Socialiste România, Editura Academiei Republicii Socialiste România.
- Schwarzová T (1967) Beitrag zur Lösung taxonomischer Probleme der *Festuca vaginata* W. et K. und *F. psammophila* Hack. Acta Fac Rer Nat Univ Comen Botanica 14: 381-414.
- Armonien R, Stukonis V, Paplauskien V (2010) Sustainable use of genetic diversity in forage and turf breeding. Sustainable Use of Genetic Diversity in Forage and Turf Breeding, pp. 41-45.
- Bednarska I, Kostikov I, Tarieiev A, Stukonis V (2017) Morphological, karyological and molecular characteristics of *Festuca arietina* Klok - A neglected psammophilous species of the *Festuca valesiaca* agg. Eastern Europe. Acta Biologica Cracoviensia Series Botanica 59(1): 35-53.
- Galli Z, Penksza K, Kiss E, Bucherna N, Heszky L (2001) Molecular taxonomic analysis of *Festuca* species: RAPD and AP-PCR analysis of the *F. ovina* group. Növénytermelés 50: 375-384.
- Galli Z, Penksza K, Kiss E, Sági L, Heszky LE (2006) Low variability of internal transcribed spacer rDNA and trnL (UAA) intron sequences of several taxa in the *Festuca ovina* aggregate (*Poaceae*). Acta Biologica Hungarica 57(1): 57-69.
- Šmarda P, Šmerda J, Knoll A, Bureš P, Danihelka J (2007) Revision of Central European taxa of *Festuca* ser. *Psammophilae* Pawlus: morphometrical, karyological and AFLP analysis. Plant Systematics and Evolution 266: 197-232.
- Šmarda P (2008) DNA ploidy level variability of some fescues (*Festuca* subg. *Festuca*, *Poaceae*) from Central and Southern Europe measured in fresh plants and herbarium specimens. Biologia 63(3): 349-367.
- Šmarda P, Kočí K (2003) Chromosome number variability in Central European members of the *Festuca ovina* and *F. pallens* groups (sect. *Festuca*). Folia Geobotanica 38(1): 65-95.
- Pawlus M (1985) Systematyka i rozmieszczenie gatunków grupy *Festuca ovina* Lw Polsce. Fragm Florist Geobot 29: 219-295.
- Borhidi A, Kevey B, Lendvai G, Seregélyes T (2012) Plant communities of Hungary. Akadémiai Kiadó.
- Penksza K (2003) *Festuca pseudovaginata*, a new species from sandy areas of the carpathian basin. Acta Botanica Hungarica 45(3-4): 365-372.
- Penksza K, Szabó G, Zimmermann Z, Lisztes Szabó Zs, Pápay G, et al. (2019) The taxonomic problems of the *Festuca vaginata* agg and their coenosystematic aspects. Georgikon for Agriculture 23(3): 63-76.

For possible submissions Click below:

Submit Article