

# Optimum $\gamma$ - Irradiation Dose for The Highest Biological Nitrogen Fixation of *Bradyrhizobium*

Alsaedi SA\*, Fadhel AS, Yousir Sh A and Abdulrazaq IB

Agricultural Research Directorate, Ministry of Science and Technology, Baghdad, Iraq



**\*Corresponding author:** Alsaedi SA, Agricultural Research Directorate, Ministry of Science and Technology, Baghdad, Iraq

**Submission:**  May 05, 2022

**Published:**  June 20, 2022

Volume 4 - Issue 1

**How to cite this article:** Alsaedi SA\*, Fadhel AS, Yousir Sh A, Abdulrazaq IB. Optimum  $\gamma$ - Irradiation Dose for The Highest Biological Nitrogen Fixation of *Bradyrhizobium*. J Biotech Biores. 4(1). JBB. 000577. 2022.

**Copyright@** Alsaedi SA, This article is distributed under the terms of the Creative Commons Attribution 4.0 International License, which permits unrestricted use and redistribution provided that the original author and source are credited.

## Abstract

This study was conducted to evaluate the possibility of increasing *Bradyrhizobium* efficiency in fixing biological nitrogen in soil. *Bradyrhizobium* isolates were obtained from cowpea field in certain Iraqi provinces. Isolates were purified, propagated and authenticated. Isolates were then subjected to a range of Gamma ray dose of irradiation of 0 to 800Gy. For all isolates number of nodules formed by all isolates increased with the increase of Irradiation dose to 400Gy after which numbers of nodules of all isolates were markedly reduced. Highest nodules numbers of all isolates were observed under 400GY irradiation dose. Under the same irradiation dose of 400GY numbers of nodules differ with different isolates being the highest under isolate No. 2.

## Introduction

BNF is a natural process in legume crops, where atmospheric dinitrogen ( $N_2$ ) is fixed into ammonia ( $NH_3$ ) in plant root nodules by a symbiotic form of *Rhizobia*, a Gram-negative *Proteobacteria*. It depends on legume genotype as well as *Rhizobia* strain, but it also reflects the interaction between crop growth and plant-available N in the soil [1].

The expanded interest in ecology has drawn attention to the fact that BNF is ecologically safe and that it's greater exploitation can reduce the use of fossil fuels and can be helpful in reforestation and in restoration of misused lands to productivity [2,3]. At present, BNF is of great practical importance because the use of nitrogenous fertilizers has resulted in unacceptable levels of water pollution (increasing concentrations of toxic nitrates in drinking water supplies) and the eutrophication of lakes and rivers [3-5]. While BNF may be almost sufficient to the needs of the organism, fertilizer is usually applied in a few large doses, up to 50% of which may be leached [3]. This wastes energy leads to serious pollution problems, particularly in water supplies. Accordingly, there has been considerable investment in improving the capacity of legume plants for fixing atmospheric nitrogen [6-8]. This includes development of complex inoculant formulations designed to improve biological nitrogen fixation, and overall plant productivity [9].

Gamma Irradiation of legume plants as a host and/or *Rhizobia* were evaluated to improve the symbiosis through the effects on nodules referred to the first steps mechanism of nodule formation, as well as, inducing re-combinations in bacterial strains through conjugation to be selecting the efficient one in symbiosis to improve nodulation process. Gamma ( $\gamma$ ) radiation has a narrow range of length high energy penetration power resulting from the nuclear disintegration of certain radioactive substances such as the isotopes Cobalt 60 ( $^{60}Co$ ) and Cesium 137 ( $^{137}Cs$ ).

Effects of radiation are accumulative [10]. Better N -fixing symbiosis may be brought by manipulating both rhizobia and plant hosts and by eventually creating an artificial rhizosphere. Numerous studies were conducted to induce genetic variations in *Bradyrhizobium* isolates to select the one of best symbiosis through alteration the expression of nodulin genes using gamma irradiation.

Therefore, this study was conducted to determine dose response curve of Gram-positive *Bradyrhizobia* isolates obtained from cowpea root in the field of Iraqi southern provinces. Gamma irradiation dose response curve of selected isolates will be utilized to determine the optimum Gamma Irradiation dose for the most efficient BNF of *Bradyrhizobium* isolate.

## Methods

### Collection of nodules

Cowpea root nodules were collected from different field sites in governorates in Iraq, belonging to the most important cowpea production area in Iraq (Basrah, Maysan and Al Anbar). Nodules were obtained and transferred into nodules collection vials as it was described by [11]. The vials were labeled and were stored at  $4\pm 1$  °C for short-term storage.

### Bacterial isolates

Nodules were surface sterilized in series of surfactants in laminar air flow cabinet by immersing in 95% v/v ethanol for 10 seconds. Followed by 30% v/v solution of sodium hypochlorite for 4min and then rinsed with sterile water. After that, they were placed in 3% v/v hydrogen peroxide followed by 5 times rinse in sterile water. The sterilized nodules were crashed with sterilized glass rod putting 1ml sterilized distilled water in petri-dish. Furthermore, the nodule suspension was streak inoculated on Yeast Mannitol Agar (YMA) [12] and incubated at 28 °C. Single colonies were selected and streaked onto YMA slant and kept at  $4\pm 1$  °C for short term storage with subculturing every 4 month. Long term storage was made by storing the culture broth in 10% glycerol at -20 °C [13].

### Culture of cowpea rhizobium on YMA-CR media

Different native cowpea rhizobium isolates were obtained from the cowpea root nodules collected from the respective sites were cultured by streak on YMA-CR medium under aseptic conditions. All the plates were labeled and incubated at 28 °C in inverted position and left for bacterial growth for 7 to 10 days in incubator; morphology of colony, nature of bacterial colony with Congo Red and shape and size of bacteria were observed for all isolates.

### Authentication of isolates by infection tests

Isolates obtained from the cowpea root nodules were authenticated as the cowpea rhizobium, the microsymbiont of cowpea. That is by examining their ability to infect cowpea roots to stimulate the root nodulation. Infection test was performed in sterile sand soil [14,15]. Sand was autoclaved for 3 times at 15 lbs (121 °C) for 20 minutes. Sterilized seeds of cowpea local cultivars were inoculated with 8 days old liquid of different bacterial isolates inocula prepared on Yeast Extract Mannitol Broth (YMB). The inoculated and uninoculated cowpea plants were grown for 4-5 weeks and examined the nodulation on their roots in both inoculated and uninoculated plants.

### Yeast manitol agar-bromothymol blue (YMA-BTB) reaction

All authenticated cowpea rhizobia cells, single bacterial colonies from the pure culture of all isolates were streaked on the plates containing YMA with 2.5ml per liter of 1.0% alcoholic bromothymol blue (YMA-BTB). Then, the plates were labeled and incubated at 28 °C. The authenticity of each isolate was detected for 10 days incubation period, with observation for color of the dye indicator on the plates [12]. Fast growing rhizobia changed color of dye indicator to yellow while slow growing rhizobia turned the dye indicator to blue.

### Preparation of liquid cowpea rhizobium inocula

Cowpea Rhizobium isolates was maintained in liquid medium i.e., Yeast Extract Mannitol Broth (YMB) for inoculation on cowpea. YMB was prepared according to Vincent [16]. Five ml of YMB was dispensed on conical flasks and sealed with aluminum foil and autoclaved. The autoclaved YMB flasks were then inoculated with cowpea rhizobium isolates, maintaining two replicates for each isolate. The cowpea rhizobium colonies from the YMA slants were transferred to the YMB flasks by sterilized inoculation loop. All activities were performed in laminar air flow cabinet. The inoculated YMB flasks were then incubated at 28 °C for 7 days in horizontal shaker with 120rpm of rotation frequency.

### Gamma irradiation of cowpea rhizobium isolates

This part of the work has been done in March 2015, at the laboratories of Soil and fertility department-IAEA / Vienna, Austria in Siebersdorf. The wild type of cowpea rhizobium isolate was inoculated into 40ml YMB until turbidity (OD600) reached 0.15 approximately ( $10^7$ CFUs/ml). The bacterial suspensions were then irradiated at 0, 50, 200, 400 and 800Gy respectively. Irradiation was done at IAEA laboratories (Seibersdorf), using a cobalt -60 source (Dose rate 2Gy/Sec.).

### Mutant/s efficiency evaluation

Soil sand mixture was Autoclaved for 30 minutes at 121 °C. Five seeds, previously one hour soaked in water, of cowpea were planted in each pot. 100mL of specified inoculant were added per pot. Each inoculant was replicated 3 times in randomized complete block design. After emergence plants were thinned to just 2 plants per pot. Phosphorus in 10mg per Kg soil sand mixture was added to each pot. Plants were grown for six weeks in a green house at  $20\pm 1$  °C day temperature and  $16\pm 1$  °C for 10 hours night.

### Preparation of bacterial cell suspension

Suspension was prepared using Yeast Extract Mannitol Broth (YMB). Five hundred milliliters of this broth were prepared and autoclaved for 20 minutes at 121 °C. Colonies from each irradiated isolates with (0, 50, 200, 400 and 800Gy) were incubated on Agar for 24hr. of these inoculants enough were transferred to Yeast Extract Mannitol Broth (YMB) and incubated for 5 days in shaker incubator

at 28 °C until growth reached the log phase at which turbidity at OD600 was 0.451. At this turbidity level there was approximately  $6.0 \times 10^8$  CFUs/ml.

### Plant harvesting

Plant root of each treatment were obtained by removing soil particle by light stream of water. Water on Plant roots was removed by soft tissue and number of nodules was calculated.

## Results and Discussion

### Authentication test

The cowpea rhizobium bacterial isolates isolated from 3 Governorates in Iraq, were tested to identify their nodulation ability. Isolated bacteria were authenticated in nodulating cowpea, their original host plant, confirming their symbiotic status.

### Bromothymol blue reaction

Bromothymol blue technique is used in bacteriology as a pH indicator in the agar, it changes to yellow in case of acid production or changes to deep blue in case of alkalization. This colorant is essentially used to differentiate Rhizobia, especially *Bradyrhizobium*

which is Alkaline in reaction compared to Rhizobium isolates which is Acidic reaction [17]. The study showed colonies reaction of all the cowpea rhizobia on Bromothymol Blue (BTB) agar plates are slow growing group (*Bradyrhizobium spp.*).

### Colony characteristics and growth response on YMA-CR medium

The bacterial colonies grown in YMA-CR plates were creamy white to translucent watery, smooth, circular and rose. The bacterial colonies did not absorb red color of Congo red when cultured in dark, which is the characteristic feature of bradyrhizobial colonies [18].

### Soils of Siebersdorf

Soils of Siebersdorf (Table 1) used in this study is sandy clay loam soil with organic matter content in the range 5-5.2%. Total nitrogen content is 2.27% with  $24.80 \text{ mgKg}^{-1}$  soil available P. According to these characteristics the soil is of good fertility level [19]. Adequate N and high plant available P will make this soil is appropriate for BNF process. The soil is also rich in Cation's content due to its high CEC.

**Table 1:** Characteristics of soil from Siebersdorf/ Austria used in the study.


Properties		Unit	Soil Texture
Separations	Clay	$\text{gmKg}^{-1}$	333
	Silt		151
	Coarse silt		106
	Fine sand		179
	Coarse sand		231
pH 1:1			7.95
Total N		%	2.27
Total P		ppm	1115
Organic matter		%	5.52
Organic Carbon		%	3.21
C/N ratio			14.14
CEC		$\text{Meq}100^{-1}$	25.38
SAR			
Ion Exchange	Ca	$\text{Meq } 100^{-1}$	22.03
	Mg		2.78
	Na		0.03
	K		0.22
	Mn		0.01
S(Ca,Mg, K, Na)			25.06
CEC			25.38
S/CEC		%	98.74

### Effect of gamma irradiation on cowpea isolates


Most efficient isolates of *Bradyrhizobium* subjected to a range of  $\gamma$ -irradiation for inducing genetic variability. Isolates were initially grown in broth then subjected to assigned dose of  $\gamma$ -radiation. Irradiated isolates were transferred to YMA media for propagation. Cowpea seeds were inoculated with each irradiated isolate.

### Effect of $\gamma$ radiation on number of nodules


Response of three isolates of *Bradyrhizobium* to given dose range of Gamma Irradiation of 0 to 800Gy was given in Figures 1-3. Isolates response was measured by the number of nodules found on plant root.



**Figure 1:** Response of *Bradyrhizobium* isolate 1 to a range of  $\gamma$  ray dose.



**Figure 2:** Response of *Bradyrhizobium* isolate 2 to a range of  $\gamma$  ray dose.



**Figure 3:** Response of *Bradyrhizobium* isolate 3 to a range of  $\gamma$  ray dose.


The data clearly showed that number of nodules formed by each isolate was increased with the increase of gamma radiation dose to 400Gy. Accordingly, Gamma dose of 400Gy is the optimum

level of Irradiation dose for the most efficient nodulation of the three isolates. At 800Gy dose, number of nodules were markedly reduced under all three isolates which may indicate that this dose

of Gamma irradiation is high enough to cause serious damage to isolate cells. These results are in agreement with those of [20-22] who had reported that Gamma Irradiation.

Figure 4 shows number of colonies formed by each isolate subjected to 400Gy dose of Gamma ray. The results clearly showed that number of nodules at the same radiation dose differ with

different Isolates. Obviously, Isolate no. 2 showed the highest number of nodules which may indicate that this isolate under the given experimental condition is the most efficient isolate. Accordingly, one may conclude that isolates nodulation efficiency increase with the increase of radiation dose to certain limit after which nodules number will decline. Furthermore, at same irradiation dose numbers of nodules differ with different isolates.



**Figure 4:** Number of Cowpea root nodules formed by 3 *Bradyrhizobium* isolate irradiated with 400Gy  $\gamma$  radiation.

## Conclusion

Methodology of using gamma irradiation technique to obtain new bacterial mutant of higher efficiency in fixing atmospheric Nitrogen was established. Gamma ray dose of 400Gy was found to be the optimum irradiation dose for the highest possible BNF efficiency of *Bradyrhizobium*.

## References

- Jensen ES, Peoples MB, Hauggaard-Nielsen H (2009) Faba bean in cropping systems. *Field Crops Research* 115(3): 203-216.
- Burris RH (1994) Biological nitrogen fixation-past and future. In: Hegazi NA, Fayez M, Monib M (Eds.), *Nitrogen fixation with nonlegumes*, The American University in Cairo Press, Cairo, Egypt, pp. 1-11.
- Sprent JI, P Sprent (1990) *Nitrogen fixing organisms. Pure and applied aspects*. Chapman & Hall, London, United Kingdom.
- Al-Sherif EM (1998) Ecological studies on the flora of some aquatic systems in Beni-Suef district. Cairo University (Beni-Suef Branch), Beni-Suef, Egypt.
- Dixon ROD, CT Wheeler (1986) *Nitrogen fixation in plants*. Thomson Science and Professional.
- Buttery BR, Park SJ (1993) Characterization of some non-fixing mutants of common bean (*Phaseolus vulgaris L.*). *Canadian J. Plant Science* 73: 977-983.
- Hardarson G (1993) Methods for enhancing symbiotic nitrogen fixation. *Plant and Soil* 152: 1-17.
- Devi MJ, Sinclair TR, Beebe S, Rao IM (2013) Comparison of common bean (*Phaseolus vulgaris L.*) genotypes for nitrogen fixation tolerance to soil drying. *Plant and Soil* 364: 29-37.
- Noel T, Parker R, Ring SG (1996) A comparative study of the dielectric relaxation behavior of glucose, maltose and their mixtures with water in the liquid and glassy states. *Carbohydrate Research* 282(2): 193-206.
- Smith-Keary PF (1975) *Mutation. Genetic structure and function*, Macmillan press LTD, Basingstoke, London, pp: 153-177.
- Al-Saedi SA, Razaq IB, Ali NA (2016) Utilization of 15N dilution analysis for measuring efficiency of biological nitrogen fixation under soil salinity stress. *Journal of Pharmacy and Pharmaceutical Sciences* 5(1): 1468-1479.
- Somasegaran P, Hoben HJ (1994) *Handbook for rhizobia. Methods in Legume-Rhizobium Technology*. Springer-Verlag, New York, USA, pp. 332-341.
- Tatiana KW, Ibrahima N, Serge B, Benoit S, Philippe de L, et al. (2003) Diversity of indigeneous bradyrhizobia associated with three cowpea cultivars (*Vigna unguiculata (L.) Walp.*) grown under limited and favorable water conditions in Senegal (West Africa). *African Journal of Biology* 2(1): 13-23.
- Namraj DR (2005) Study on effectiveness of native bradyrhizobial isolates for N2-fixation in soybean (*Glycine max (L.) MERR.*). Central Department of Botany Tribhuvan University Kathmandu, Nepal.
- Berns AE, Philipp H, Narres HD, Burauel P, Vereecken H, et al. (2008) Effect of gamma-sterilization and autoclaving on soil organic matter structure as studied by solid state NMR, UV and fluorescence spectroscopy. *European Journal of Soil Science* 59(3): 540-55.
- Vincent JM (1970) *A manual for the practical study of the root nodule bacteria*. International Biological Program, Blackwell Scientific Publications, New Jersey, USA.
- Jacinta K, Muthuri (2014) Symbiotic effectiveness of *bradyrhizobium japonicum* USDA 110 and *sinorhizobium FREDII* USDA 191 on two different soybean cultiv ARS. *European Scientific Journal* 10(12): 329-337.
- Usharani G, Jayanthi M, Kanchana D, Saranraj P (2014) Effect of *bradyrhizobium* isolates for the maximization of growth and yield of black gram (*Vigna mungo L.*). *International Journal of Advance Multidisciplinary Research* 1(4): 33-39.
- Kim HT (2011) *Principles of soil chemistry*. (4<sup>th</sup> edn), pp. 210-220.

20. Min J, Lee CW, Gu MB (2003) Gamma radiation dose-rate effects on DNA damage and toxicity in bacterial cells. *Radiat Environ Biophys* 42(3): 189-192.
21. Vilenchik MM, Knudson AG (2006) Radiation dose-rate effects, endogenous DNA damage, and signaling resonance. *Proc Natl Acad Sci USA* 103(47): 17874-17879.
22. Chitchanok A, Rattasarith P, Suthatip S, Suphaporn P, Nattayana P, et al. (2011) Improvement of vitamin B6 production from rhizobium sp. 6-1C1 by random mutation. *KKU Res J* 16(8): 911-918.

For possible submissions Click below:

[Submit Article](#)