



Breast Cancer Chemotherapy and Treatments



Li Pin Kao^{1,2*}

¹Department of Basic Medical Sciences, Purdue University, Indiana, USA

²Centre for Cancer Research, Purdue University, Indiana, USA

*Corresponding author: Li Pin Kao, Department of Basic Medical Sciences, Centre for Cancer Research, Purdue University, USA

Submission: February 02, 2018; Published: August 17, 2018

Abstract

Breast cancer, the first most common malignant tumor in women worldwide. Many etiological factors such as a wide spectrum of clinical manifestations caused by family clustering, hormonal factors, physiology changed, environmental inducers, and life styles. Although there are many FDA-approval drugs for breast cancers. Tumors are heterogenous and individual has inherited different genetic background. Thus, it is hard to find an effective treatment and it may recurrent after recovery from first treatment.

Keywords: Breast cancer; Autophagy; Autophagy related inhibitors; Chemotherapy

Introduction

The mortality rate exceeds 144,000 cases and most of the breast cancer (BC) cases are resistant to traditional chemotherapy and radiotherapy. Wide varieties of chemotherapeutic agents have been tried and are in use, including tamoxifen, docetaxel, anthracyclines (i.e. doxorubicin, and epirubicin), taxanes (i.e. paclitaxel, and docetaxel), 5-fluorouracil, cyclophosphamide, and carboplatin. No regimen has been proved to be curative. Often, the response rate and prolongation of survival are minimal (a few months or less), and there is a significant morbidity associated with poor treatment effects. Surgical resection is considered the first-line options for early tumors, although there is no agreement on which the best approach is. Interestingly, triple negative has been reported has higher chemo-resistance than other type of breast cancers with low autophagy activity [1,2]. Antiestrogen resistant cell lines exhibit increased basal autophagy when compared with their antiestrogen sensitive parental cells [3]. Study has shown to manipulate transient receptor potential channel 5 (TRPC5), a Ca₂₊ permeable cation channel, which helps in promotes autophagic activity [4]. Studies also demonstrate the breast cancer usually Prosurvival after autophagy activity increase by various therapeutics [5-7]. Moreover, Tamoxifen and Faslodex (ICI) both induce autophagy in ER+ breast cancer cells without [3-11]. There are small molecules

Table 1: Autophagy inhibitors during autophagy pathway.

which has been in FDA-approval drug lists and applied in clinical trial in different diseases.

Antiestrogen resistant cell lines exhibit increased basal autophagy. Study had demonstrated combination of Tamoxifen or Faslodex (ICI) with Hydroxychloroquine (HCQ) had different anti-estrogen responsiveness *in vitro* or *in vivo* which may affect by tumor microenvironment (i.e. chemokines, macrophage development/activity [12]. Inhibiting autophagy via autophagy-related genes (i.e. autophagy-related gene (Atg) 5, Atg7, and p62/SQSTM1) silencing potentiated antiestrogen-mediated cell death, indicating that antiestrogen stimulated autophagy is pro-survival and a critical mechanism of therapy resistance. Overall, it indicated that increased in autophagy activity in early recurring breast cancer when compared with breast cancer that never recurs. Moreover, elevated p62 is significantly correlated with poor survival in breast cancer patients, suggesting a role for autophagy in breast cancer re-occurrence. Manipulation of autophagy activity can be a potential therapy for chemotherapy *in vitro* or *in vivo*. Therefore, diagnosis or detection at an early stage is crucial to allow the application of treatments for increasing the life expectancy of the patient [13] (Table 1,2).

Name	Mechanism	
3-Methyladenine	phosphoinositide3-kinase (PI3) inhibitor	Auto phagosome formation
Wortmannin	PI3-kinase inhibitor	Autophagosome formation
LY294002	PI3-kinase inhibitor	Autophagosome formation
SBI-0206965	Unc-51-like kinase 1 (ULK1) Inhibitor	Autophagosome formation

Spautin-1	ubiquitin-specific peptidases (USP10) and (USP13) inhibitor	Autophagosome formation
SAR405	Vacuolar Protein Sorting Protein 18 and 34 (Vps18 and Vps34) inhibitor	Autophagosome formation
NSC185058	autophagy-related gene 4 (ATG4) inhibitor	Autophagosome formation
Verteporfin	Unknown	Autophagosome formation and accumulation
ROC325	Unknown	Lysosome
Lys05	Unknown	Lysosome
Chloroquine	Unknown	Lysosome
Hydroxychloroquine	Unknown	Lysosome

Table 2: Current Hydroxychloroquine (HCQ) clinical trials.

Treatment	Condition	Phase Trial	Reference # at ClinicalTrials.gov
HCQ + sunitinib malate	Adult solid neoplasm	I	NCT00813423
HCQ + vorinostat	Malignant solid tumour	I	NCT01023737
HCQ + sirolimus or vorinostat	Advanced cancers	I	NCT01266057
HCQ + Protein kinase B (Akt) inhibitor MK-2206 dihydrochloride (MK2206)	Advanced cancers	I	NCT01480154
HCQ as a single agent	Estrogen receptor positive breast cancer	I	NCT02414776
HCQ + gemcitabine	Advanced adenocarcinoma	I/II	NCT01506973
HCQ + Interleukin 2(IL-2)	Renal cell carcinoma	I/II	NCT01550367
HCQ + vorinostat	Colorectal cancer	I/II	NCT02316340
HCQ + gemcitabine/carboplatin	Small cell lung cancer	I/II	NCT02722369
HCQ + capecitabine	Pancreatic carcinoma	II	NCT01494155
HCQ as a single agent	Prostate cancer	II	NCT00726596
HCQ + Abraxane and gemcitabine	Pancreatic carcinoma	II	NCT01978184

Conclusion

It is very little is known how the manipulation of a biological process toward to potential cancer therapy. As is often the case, we need understand the basic mechanistic of biological processes if this is going to apply for a clinical benefit or other practical application. This article demonstrated the importance of manipulation of autophagy activities in different cancer treatment in basic biomedical research and different clinical trials. Given the rapid progress in understanding autophagy in cancer-related researches and what it does, manipulation of autophagy activities in cancer-related research has ready to take off.

Acknowledgement

The author would like to acknowledge the financial support of the Purdue University for provided support in the form of salaries of KLP listed but did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- Garbar C (2017) Chemotherapy treatment induces an increase of autophagy in the luminal breast cancer cell MCF7 but not in the triple-negative MDA MB231. *Sci Rep* 7(1): 7201.
- O'Reilly EA (2015) The fate of chemo resistance in triple negative breast cancer (TNBC). *BBA Clin* 3: 257-275.
- Cook KL (2012) Glucose regulated protein 78 controls cross talk between apoptosis and autophagy to determine antiestrogen responsiveness. *Cancer Res* 72(13): 3337-3349.
- Zhang P (2017) TRPC5 induced autophagy promotes drug resistance in breast carcinoma via CaMKKbeta/AMP alpha/mTOR pathway. *Sci Rep* 7(1): 3158.
- Clarke R (2012) Endoplasmic reticulum stress the unfolded protein response autophagy and the integrated regulation of breast cancer cell fate. *Cancer Res* 72(6): 1321-1331.
- Thomas S (2011) Addition of a histone deacetylase inhibitor redirects tamoxifen treated breast cancer cells into apoptosis which is opposed by the induction of autophagy. *Breast Cancer Res Treat* 130(2): 437-447.
- Vazquez Martin A, C Oliveras Ferraros, JA Menendez (2009) Autophagy facilitates the development of breast cancer resistance to the anti HER2 monoclonal antibody trastuzumab. *PLoS One* 4(7): e6251.
- Cook KL (2014) Mitochondria directly donate their membrane to form autophagosomes during a novel mechanism of parkin associated mitophagy. *Cell Biosci* 4(1): 16.
- Clarke R (2009) Gene network signalling in hormone responsiveness modifies apoptosis and autophagy in breast cancer cells. *J Steroid Biochem Mol Biol* 114(1-2): 8-20.
- Schoenlein PV (2009) Autophagy facilitates the progression of ER alpha positive breast cancer cells to antiestrogen resistance. *Autophagy* 5(3): 400-403.
- Samaddar JS (2008) A role for macroautophagy in protection against 4-hydroxytamoxifen-induced cell death and the development of antiestrogen resistance. *Mol Cancer Ther* 7(9): 2977-2987.

12. Cook KL (2014) Hydroxychloroquine inhibits autophagy to potentiate antiestrogen responsiveness in ER+ breast cancer. Clin Cancer Res 20(12): 3222-3232.


13. Chude CI, RK Amaravadi (2017) Targeting Autophagy in Cancer: Update on Clinical Trials and Novel Inhibitors. Int J Mol Sci 18(6).



Creative Commons Attribution 4.0 International License

For possible submissions Click Here

[Submit Article](#)



Modern Approaches in Drug Designing

Benefits of Publishing with us

- High-level peer review and editorial services
- Freely accessible online immediately upon publication
- Authors retain the copyright to their work
- Licensing it under a Creative Commons license
- Visibility through different online platforms