



# Climate Change and Sustainable Management of Salinity in Agriculture



Gurbachan Singh\*

Ex-Chairman of Agricultural Scientists Recruitment Board, India

\*Corresponding author: Gurbachan Singh, Beant Villa Building, Adjacent Adarsh Public School, Near Power House, Kunjpura Road, Karnal-132001, Haryana, India, Email: dr.gurbachan@gmail.com

Submission: June 15, 2018; Published: August 09, 2018

## Introduction

There has been a steady increase in the emission of greenhouse gases like carbon dioxide, methane and nitrous oxide in the environment. Agriculture is reported responsible for up to almost half of all methane emissions. The climate change predictions over India indicate that temperature rise is likely to be around 3°C and rainfall increase is expected by 10-20 per cent over central states of India by 2100 A.D. The climate change triggered frequency of weather related events like floods, droughts, frost, cold and heat waves has considerably increased during last two decades. Continuation of such trends associated with rise in temperature is expected to melt ice, glaciers, re-distribute water flow in rivers, raise sea levels, sub-merge coastal habitats, islands, generate tsunamis and dislocate human and livestock settlements. Predicted spatial redistribution of precipitation, droughts, floods and water balance will change land use, pests, diseases and other ecological parameters. These changes will necessitate the need to devise research strategies to deal with predicted changes to sustain agricultural productivity and to achieve food and nutritional security in 21<sup>st</sup> century.

Soil salinity and sodicity has degraded about 6.73 million ha otherwise productive area in the country. Development and refinement of appropriate technologies for reclamation and management of such lands seems promising option to achieve future food and nutritional security as well as adoption and mitigation strategy to cope with predicted future climate change scenario. In the present paper, information has been compiled and discussed how reclamation and management of salt - affected soils and waters will help sustaining food security and also help moderating/negating climate change related risks in the near future through carbon sequestration in vegetation and soil. The information has been compiled under the sub heads: future climate change scenario in India, climate change and agriculture Vulnerability, impact of climate change on agriculture: recent case studies, harnessing salt-affected soils potential and future research and development needs.

## Future climate change scenario in India

Table 1: Climate change projections for India.

Year	Sea-son	Temperature Change (OC)		Rainfall Change (%)	
		Lowest	Highest	Lowest	Highest
2020s	Annual	1.00	1.41	2.16	5.97
	Rabi	1.08	1.54	-1.95	4.36
	Kharif	0.87	1.17	1.81	5.10
2050s	Annual	2.23	2.87	5.36	9.34
	Rabi	2.54	3.18	-9.22	3.82
	Kharif	1.81	2.37	7.18	10.52
2080s	Annual	3.53	5.55	7.48	9.90
	Rabi	4.14	6.31	-24.83	-4.50
	Kharif	2.91	4.62	10.10	15.18

Source: Lal [1]

The central research institute for dryland agriculture through its all India coordinated research projects on agro-meteorology and dryland agriculture has worked out the trends in maximum and minimum temperature during Kharif and Rabi season for Indian conditions. Their calculations indicate that there may be several location specific uncertainties in the minimum and maximum temperatures which may have adverse impact on agricultural production and productivity. Their calculations for trends in monthly rainfall in various sub-divisions in India indicate that rainfall may be more during December, January and February in case of West Rajasthan, Punjab and Haryana. However, there may be slight decrease in October and November rainfall in several divisions. The analysis was also made about trends of rainfall in India and shift in surplus rainfall for the period 1871 to 1999 and 1960 to 1999. The observations indicate a shift in surplus rainfall from west towards east. Lal [1] made calculations about the

predicted climate change and its impact on agriculture in India and the results are reported in Table 1. He reported an annual mean area averaged surface warming over Indian Subcontinent to range between 3.5 to 5.5 °C over the region by 2080. These projections showed more warming in winter season over summer monsoon. Rise in surface temperature in north India is predicted 3 °C or more by 2050. A marginal increase of 7 to 10% in rainfall is predicted over the subcontinent by 2080. Further, the study revealed a fall in rainfall by 5 to 25% in winter months and an increase of 10 to 15% in summer monsoon rainfall in India. These projected changes will have both beneficial and adverse effects on agriculture, horticulture, environment and socio-economic set-up. To devise strategies to manage such changes and to reduce vulnerability of agriculture, careful analysis of old climate data and linking with weather forecasting and working out eco-years, eco-seasons, eco-months and eco-weeks at each agricultural university and ICAR research institute will be required. Samra and Singh (2002) suggested several strategies and contingent crop plans to negate/moderate the impact of sub-due rainfall/drought in different agro-meteorological sub divisions of the country.

It has been estimated that mountain glaciers in the Himalayas and on the Tibet-Qinghai Plateau are melting and could deprive the major rivers of India and China of the ice melt needed to sustain them during the dry season in the future. Scientists believe that water flows in Yellow river and the Yangtze River basins where irrigated agriculture depends heavily on river will decrease and may have negative impact on food production. However, the scientists at the National Institute of Hydrology (NIH, Roorkee) have studied the glaciers in the Himalayas and their results indicate that situation is not that serious but there has been some decrease in the size of glaciers at some locations. There is a strong need to bench-mark the melting processes and their annual monitoring using GPS and satellite systems.

### Salinity problem: global and Indian scenario

Soil and vegetation represent potential sinks for carbon sequestration and reforestation is considered as a possible means of mitigation of global warming. As per recent estimates nearly 953 million ha area is affected by salinity and sodicity in the world (Table 2). Australia followed by north and central Asia and south America have the maximum problem of high salt concentration in the root zone soil. State wise area affected by salinity and sodicity in India is given in Table 3. As per reconciled data reported in Table 3, about 6.73 million ha otherwise productive area is constituted by salty lands. Gujarat state followed by UP has maximum area affected by salinity and sodicity. It has been further projected that 11.7 million ha area will be affected by 2025.

**Table 2:** Distribution of salt affected soils in the world (million ha).

Region	Soionchaksl Saline	Solonetzl Sodie	Total
North America	6	10	16
Mexico and Central America	2	-	2

South America	69	60	129
Africa	54	27	81
South and West Asia	83	2	85
South and East Asia	20	-	20
North and Central Asia	92	120	212
Australia	17	340	357
Europe	-	-	51
Total	-	-	953

**Table 3:** Extent of salt affected soils in India (ha).

State	Saline	Sodic	Total
Andhra Pradesh	77598	196609	274207
Andaman & Nicobar Island	77000	0	77000
Bihar	47301	105852	153153
Gujarat	1680570	541430	2222000
Haryana	49157	183399	232556
Karnataka	1893	148136	150029
Kerala	20000	0	20000
Madhya Pradesh	0	139720	139720
Maharashtra	184089	422670	606759
Orissa	147138	0	147138
Punjab	0	151717	151717
Rajasthan	195571	179371	374942
Tamil Nadu	13231	354784	368015
Uttar Pradesh	21989	1346971	1368960
West Bengal	441272	0	441272
Total	2956809	3770659	6727468

**Table 4:** Percentage use of poor quality waters in different states.

State	Percentage (estimated values)
Andhra Pradesh	32
Gujarat	30
Haryana	62
Karnataka	38
Madhya Pradesh	25
Rajasthan	84
Uttar Pradesh	47

Source: CSSRI, Karnal

Several studies in India [2] reveal that extremely carbon depleted soils like salt-affected soils have quite high potential for sequestering carbon in vegetation and soil if suitable tree and grass species are grown along with best management practices like rain water conservation. Nearly 25% of the ground water resources in India are saline and/or brackish. Continuous use of such water for irrigation is bound to increase soil salinity/sodicity. States like Rajasthan and Haryana have 84% and 62%, respectively ground

water as saline/sodie. Percentage use of poor quality waters in selected states is given in Table 4.

Introduction of irrigation in canal command areas coupled with over use of water has resulted in rise in ground water. In some of the canal commands water table is rising @ 0.3 to 1.2 meter/annum (Table 5). Nearly 3.55 million ha is already water logged and gone out of cultivation. Statewise extent of water logged area is reported in Table 6. Losses in productivity due to water logging are reported 42% in paddy, 38% in wheat, 77% in cotton and 61% in sugarcane compared to productivity of these crops in normal soils (Table 7).

**Table 5:** Rising trend of water table in irrigation commands.

Irrigation Command	Rise of Water Table (Meter/Annu)
IGNp, Rajasthan	0.29 - 0.88
Western Yamuna and Bhakra canal	0.30 - 1.00
Sharda Sahayak, U.P.	0.68
Nagarjuna Sagar, A.P.	0.32
Malprabha canal command, Karnataka	0.60 - 1.20

**Table 6:** Extent of water logged areas in India.

State	Waterlogged Area (mha)
Andhra Pradesh	0.43
Bihar	0.35
Gujarat	0.17
Haryana	0.23
Karnataka	0.05
Kerala	0.08
Madhya Pradesh	0.06
Maharashtra	0.02
Orissa	0.18
Punjab	0.30
Rajasthan	0.18
Tamil Nadu & Pondicherry	0.61
Uttar Pradesh	0.59
West Bengal	0.29
Total	3.55

Source: Ministry of Agriculture (Draft Report, 1999)

**Table 7:** Losses due to water logging and soli salinity.

Crop	Normal Lands	Salt Affected Lands	Waterlogged Lands
Paddy	39.9	21.8(45)	23.0(42)
Wheat	26.0	15.8 (40)	18.6 (38)
Cotton	16.3	6.1 (63)	3.7 (77)
Sugarcane	636.8	330.2 (48)	247.5 (61)

### Reclamation and management strategies

After the establishment of Central Soil Salinity Research Institute in 1969 at Kamal, several site-specific technologies have

been developed for reclamation and management of saline and alkali soils. A gist of the relevant technological options is discussed as under: Sub-surface drainage technology has been developed to reclaim waterlogged saline soils. State-wise drained area covered is given in Table 8. Similarly, change in cropping intensity after installation of the drainage system is reported in Table 9.

**Table 8:** state wise drainage area covered.

State	Irrigation Command	Area Covered (ha)
Rajasthan	Chambal	15,700
	Indira Gandhi Nahar Pariyojna	500
Haryana	Eastern Jamuna Canal	1450+3000
	Bhakhra Canal	1300+1000
Punjab	South West Punjab	30+2000
Karnataka	Upper Krishna	30
	Tungabhadra	200
	Malparhba Ghatparbha	20
Andhra Pradesh	Nagarjuna Sagar	50
	Krishna Western Delta	50
Madhya Pradesh	Unspecified	50
Maharashtra	Uncommanded/ Neera Canal	1000+1000
	Command/others	
Gujarat	Mahi-Kadana	150
	Ukai- Kakrapar	80
Kerala	Acid Sulphate Soils	30
Assam	Tea Gardens	15

**Table 9:** Change in cropping intensity.

Place	Before Drainage	After Drainage	Increase (%)
Sampla	0	200	-
Ismaila	73	148	103
Gohana	117	175	50
Konanaki	70	90	29
Uppugunduru	130	165	27
Islampur (Karnatka)			
ORP	0	200	-
Phase II	88	156	77

Several salt tolerant varieties of rice, wheat and mustard have been developed which can be grown either without or with half the recommended dose of amendments. The promising varieties are listed below:

- Rice: CSR 10, CSR 13, CSR 23, CSR 27, CSR 30 and CSR 36  
 Wheat: KRL 1-4, KRL 19, KRL210 and KRL213  
 Mustard: CS 52, CS 54 and CS 56

Besides these, salt tolerant trees, grasses, bushes, biofuel crops have been identified and agro-techniques for their successful

cultivation in salty lands standardized. Several multipurpose tree based silvi-agri and agroforestry models have been developed for fuelwood and forage production and bio-amelioration of both sodic and saline soils. An auger hole technique for establishing tree plantations in soils of  $pH_2 > 10.0$  was developed. Several village community lands affected by salinity/ sodicity have been planted with trees like *Prosopis juliflora*, *Acacia nilotica*, *Eucalyptus tereticornis*, *Casuarina equisetifolia* and *Tamarix articulata*. The augerhole technique was further improved as pit - augerhole technique which made it possible to grow even fruit trees such as almas, guava, jamun, karaunda, bear and imly in salty lands. Trees and grasses once established in salt lands help reclamation of such soils by root activity and addition of leaf litter (Table 10).

**Table 10:** Reclaiming effects of *Prosopis juliflora* plantations on village community lands.

Village	Depth (m)	Soil Properties			
		pH		Organic Carbon (%)	
		Initial	After 8 years	Initial	After 8 years
Shera	0-15	10.0	8.8	0.13	0.048
	15-30	10.1	9.6	0.06	0.21
Sutana	0-15	10.2	8.5	0.20	DAD
	15-30	10.6	9.4	0.07	0.09
Nain	0-15	9.7	8.6	0.08	0.45
	15-30	10.2	9.2	0.10	0.12
Shalsi	0-15	10.0	8.8	0.22	0.65
	15-30	10.5	10.1	0.08	0.12

Association of salt-tolerant grasses like *Leptochloa fusca* with trees further hastens the reclamation process. The short, medium and long-term effects of trees on soil properties in general and carbon sequestration in particular is reported in Table 11 & 12. The perusal of the data clearly indicates that significant amount of carbon is sequestered in the soil. The extent of carbon build-up depends upon the nature of tree species, planting density, age of plantation and the management practices.

**Table 13:** Uses of *Opuntia* species.

Food	Fruits and fruit peel, juice, pulp, alcoholic beverages, jam, syrup
Forage	stems/cladodes, fruits, seeds, cultivated as forage shrub
Energy	Biogas, ethanol, firewood
Medicine	Diarrhoea (stem), diuretic (flower, root), amoebic dysentery (flower), diabetes (stem), hyperlipidemy (stem), obesity (fibres), antiinflammatory (stem)
Cosmetics	Shampoo, cream, soaps, body lotions
Agronomic	Hedges and fences, mulching, soil improver, wind break, organic manure
Others	Adhesives and glues, pectin, fibres for handicrafts, paper (stem), Dyes (fruit), rearing of <i>Oaetyloptus oews</i> on cladodes, antitranspirant, ornamental

**Table 11:** Ameliorating effects of mesquite and other tree plantations on an alkali soil.

Species	Original		After 20 years	
	$pH_2$	Organic Carbon (%)	$pH_2$	Organic Carbon (%)
<i>Eucalyptus tereticornis</i>	10.3	0.12	9.18	0.33
<i>Acado nilotica</i>	10.3	0.12	9.03	0.55
<i>Albizia lebbek</i>	10.3	0.12	8.67	0.47
<i>Terminalia arjuna</i>	10.3	0.12	8.15	0.47
<i>Prosopis juliflora</i>	10.3	0.12	8.03	0.58

**Table 12:** Straw and grain yields of wheat and oats in soils taken under the 25 years old tree plantations and in the adjacent crop land soil.

Tree Plantations	Wheat		Oats	
	Grain (9 pot')	Straw (9 pot')	Grain (g por')	Straw (g pot')
	<i>Prosopis juliflora</i>	61.7	87.5	87.9
<i>Terminalia arjuna</i>	44.0	38.5	45.8	62.8
<i>Eucalyptus tereticornis</i>	32.2	25.3	42.7	58.5
<i>Albizia lebbek</i>	45.3	43.5	52.8	66.9
<i>Acacia nilotica</i>	55.7	68.8	61.6	67.5
Cropland	13.3	15.4	24.3	26.7
LSD (p=0.05)	2.8	2.0	7.0	9.4

Mechanism of soil reclamation by trees and grasses and processes involved are explained below in Table 8 & 9. The crops grown in the soils reclaimed by trees and trees plus grass mixtures are reported to have much better productivity compared to their productivity in soils reclaimed by amendments (Table 13).

In general, it is difficult to build organic carbon in soils where summer temperature exceeds 40-45 °C. However, integration of trees with crops in a unified agroforestry system helped building appreciable quantity of organic matter in the soil (Table 14).

**Table 14:** Change in soil properties (0-30 cm) under different tree-crop combinations in 5 years.

Land use Systems	Organic Carbon (%)	Available N (kg ha <sup>-1</sup> )
Crop based system	+0.07	+10
Eucalyptus based	+ 0.12	+21
Acacia based	+0.20	+ 31
Populus based	+0.17	+ 25

Role of trees having high transpiration rates has been exploited to lower the water table in areas where ground water is showing rising trends. This process is known as bio-drainage. Bio-drainage is defined as the process of removing the excess soil water through transpiration using bio-energy of the plant and radiation energy of the Sun. It is an option to prevent the development of water logged and saline soils especially in land locked areas where there is no possibility of disposing saline drainage effluent. Also, there are large areas which get water stagnation due to seepage from higher elevation, surface disposal of urban and industrial effluents and floods during rains, where due to their topographical locations, conventional system of de-watering through mechanical pumping and surface drainage is not possible. Promising tree species for bio-drainage use include: *Eucalyptus*, *jamun*, *shisham*, bamboo, poplar etc.

### Salt affected grazing lands

Salt affected grazing lands are predominant landscapes throughout the country. The biomass production capacity of these lands is relatively low because of poor soil health and management. The above ground biomass of saline communities in peripheral areas of runs in Rajasthan was reported ranging from 0.6 to 0.5 t ha<sup>-1</sup> while in saline depressions of Jaisalmer, some grass communities were found to produce from 0.30 to 0.65 t ha<sup>-1</sup> forage biomass. Major species which contribute significant biomass as fodder in such lands include: *Aeluropus*, *Atriplex*, *Bothriochloa*, *Cenchrus*, *Chenopodium*, *Chloris*, *Cynodon*, *Dactyloctenium*, *Dichanthium*, *Eragrostis*, *Leptochloa*, *Kochio*, *Panicum*, *Salsola*, *Suaeda*, *Tribulus*, *Salvadora*, *Prosopis* and *Ziziphus*. Similarly, the important top feed shrubs and trees include *Ailanthus excelso*, *Acacia nilotica*, *A. catechu*, *A. eucophloea*, *A. tortilis*, *Balanites roxburghii*, *Prosopis cineraria*, *P. juliflora*, *Azadirachta indica*, *Albizia julibrissin*, *Leucaena leucocephala*, *Dahlbergian sissoo*, *Melia azadirach*, *Hordwickia binata*, *Grewia ovata*, *Ficus bengalensis*, *F. religiosa*, *Anogeissus pendula*, *Bauhinia variegata*, *B. racemosa*, *Butea monosperma*, *Cordia dichotoma*, *Flacourtia indica*, *Moringa oleifera*, *Dichrostachys nuttiness*, *Morus alba*, *Ziziphus mauritiana* and *Z. nummularia*. The leaves of most of these trees are rich in nutrients. This type of fodder becomes more relevant during drought scarcity period.

### Salt tolerant halophytic pastures

Many halophytes grow in natural saline or other arid environments, which are rapidly deteriorating through the process of desertification. It is necessary for plants growing under saline conditions to maintain a high concentration of osmotic substances to successfully overcome the water-relating forces of the surrounding soil. Halophytes growing in saline soil are able to change the osmotic potential of their cell sap whenever required.

The value of certain salt-tolerant shrub and grass species have been recognized by their incorporation in pasture-improvement programmes in many salt-affected dry regions throughout the world. There have been recent advances in selecting species with high biomass and protein levels in combination with their ability to survive a wide range of environmental conditions, including salinity and drought. Considerable success has been achieved in Indian Subcontinent and elsewhere in cultivating halophytic forages such as chenopads, especially *Atriplex* in areas subject to total summer drought on badly salt-affected lands. Grazing animals can meet their requirement of protein and metabolisable energy through mixture of salt land fodders, tops of plantations of halophytic shrubs/trees (providing crude protein) and a minimal understorey cover of forages. Another plant, which has attracted much attention as a fodder plant on soils affected by salinity and waterlogging is kallar grass (*Leptochloa fusca*). In India and Pakistan, this grass has been widely utilized for production in both alkali and saline waterlogged soils. In a silvi-pastoral system with trees such as *Acacia nilotica*, *Prosopis juliflora* (the loppings of these could be used as forage) for initial four years. On an average *Leptochloa* could produce 15.6 to 17.4 t ha<sup>-1</sup> green forage per annum on highly alkaline soil (pH>10) even during low rainfall years. Species of halophytic forages like *Sarcocolla*, *Chenopodium*, *Kochia*, *Atriplex*, *Salsola*, *Suaeda*, *Trianthema*, *Portulaca*, *Tribulus* and *Alhagi* along with several grasses such as *Aeluropus/agopoides*, *Sporobolus* (*S. marginatus*, *S. airoides*, *S. diander*, *S. hexostachyus*, *S. tramuus*), *Cynodon dactylon*, *Dactyloctenium aegyptium*, *Paspalum vaginatum*, *Chloris gayana*, *Echinochloa turnerana*, *Eragrostis tenella*, *Dichanthium annulatum*, *Brachiaria mutica*, *Bathriochloa pertusa* and many others are commonly found grown naturally in both alkali and saline soils and contribute as forage even during drought period.

In many coastal areas (including low rainfall areas of United Arab Emirates and Rann of Kutch in India) where mangroves occur sporadically and there is scarcity of fodder, the foliage of many mangroves and associated plants such as species of *Pongamia*, *Barringtonia* and many others, are used as forage for cattle, goats and camel. Among other trees particularly grown on inland salty lands, species of *Acacia*, *Prosopis*, *Salsola*, *Cordia*, *Ailanthus*, *Balanites* and *Ziziphus* are traditional fodder plants of drought prone arid regions. Many species of *Acacia* (*A. cyclopes*, *bivenosa*, *implys*, *halosericea*, *saligna*, *saicina*, and *victoria*) and *Prosopis* (*P. juliflora*, *cineraria*, *chilensis*, *gianduosa*, *pauciflora* and *tamarugo*) are among the most promising genetic resources to be utilized in developing fodder sources in drought prone areas.

## Trees and bushes on field boundaries and bunds

Painting of multipurpose tree species (MPTS) and promising bushes on the boundaries of agricultural fields in drought prone salt affected area is useful as (i) biefence protection against wild animals, (ii) as an alternate source of fodder, fuel and income generating products during severe drought, (iii) as vegetative barriers to conserve soil and water, (iv) moderating effects of drought through moderating micro-climate and (v) in some cases act as shelterbelts/wind breaks and carbon sequestration.

Farmers make high bunds on the periphery on their agricultural fields, plant *Saccharum munja* on top and fix thorny bushes/branches of trees to protect their crops from wild animals. Such bio-fencing practices need replacements at shorter intervals and do not generate any product of economic value. The list of species which can be exploited as a live fence and drought moderating option includes: *Capporis decidua*, *Opuntia ficu -indica*, *Leucaena leucocephala*, *Carrisa carandas*, *Lantana camara*, *Aloe Barbadians*, *Prosopis juliflora*, etc.

## Underutilized agro forestry resources

Large number of trees, bushes, shrubs and grasses are naturally growing as wild plants in rainfed regions of the world. Such plants are adapted to rainfed situations and have tremendous potential to be exploited as a food, forage, fuel and/or industrial crops [3,4]. Some of these plants are already exploited in different parts of the world. For example, edible cactus (*Opuntia ficus indica*) is being extensively used as a fruit, forage, vegetable and medicinal crop in countries like Mexico, Argentina, France, Brazil, Italy, South Africa and even in south western USA. The other uses of cacti in arid and semi-arid areas include: live fence, vegetative barrier for soil and water conservation in sloping land and as a wind break. A summary of *Opuntia* uses is presented in Table 13. The forage quality parameters of this underutilized plant as reported in literature are cited in Table 15.

**Table 15:** Range in values for cactus cladode composition for use in animal feed.

Moisture content (%)	85-90
Crude protein (%)	5-12
<i>In vitro</i> dry matter digestibility (%)	75
<i>In vitro</i> protein digestivity (%)	72
Crude fibre (%)	43
P (0/0)	0.08-0.018
Ca (%)	4.2
K (%)	2.3
Mg (%)	1.4
Energy (Meal kg <sup>2</sup> )	2.6
Carotenoids (mg 100 g <sup>2</sup> )	29

Similarly, trees of the genus *Prosopis* are highly tolerant to aridity and salts. *Prosopis juliflora* has the potential to grow greener even during severest drought period. The trees provide livelihood support in terms of firewood, fodder and even pods as human

and cattle feed. It has also been exploited economically as timber, firewood and forage tree in countries like Brazil, Argentina, Peru, Mexico, Senegal, South Africa and USA [5-7]. Comparison of protein content of *Prosopis juliflora* in relation to other important trees as reported in the literature is given in Table 16. The data indicates that the *Prosopis* pods and leaves contain almost similar protein as that found in alfalfa.

**Table 16:** Comparison of protein content of mesquite (*Prosopis juliflora*) products in relation to other products in Brazil.

Product	Protein (%)
Pods ( <i>Prosopis juliflora</i> )	12.9
Leaves ( <i>Prosopis juliflora</i> )	13.6
Maize ( <i>Zea mays</i> )	6
Alfalfa ( <i>Medicago sativa</i> )	14.1
Guinea grass ( <i>Panicum maximum</i> )	2.6

To exploit the use of under - utilized plant resources as climate change adoption and mitigation strategy we need to focus on : (a) to prepare an inventory of under - utilized/un-exploited plants and maintain their germplasm for productivity and quality improvement, (b) identification of ecological limits and optimal management practices for promising species and their evaluation as mono-cultural mixed communities or even as under - storey crop with trees, and (c) linking research and developmental issues of under - utilized plants with already existing international networks on such plants and (d) standardization of post harvest processing techniques for value addition and marketing opportunities.

## Shelter belt

Well-designed shelterbelts conserve moisture and soil against wind erosion, neutralize heat wave, cold wave and fruit fall due to wind blowing. In addition to these effects, shelterbelts can also provide a wide range of useful products, from poles and fuelwood to fruits, fodder, fibre and mulch. They have been planted quite extensively along Indira Gandhi Rajasthan Canal, roads, railway lines and around the orchards. Species of *Eucalyptus*, *Populus*, *Cosuarina*, *Prosopis*, *Acacia*, *Leucaena*, *Azadirachta*, *Moringa* and *Gliricidia* are most frequently grown as wind breaks/shelterbelts. These are distinguished from boundary plantations and living fences by their orientation, which must face the wind. Diversifying the species in the shelterbelt can also bring a wider variety of useful products to local users.

## Conclusion

Development and refinement of technologies for rehabilitation of 120 million ha degraded/wastelands in general and about 7 million ha salt-affected soils in particular need to be considered on high priority. These carbon depleted soils will provide much needed opportunity to sequester carbon in the soil. Sequestration of carbon will not only imply increasing the amount of carbon entering the soil but also a decline in the amount leaving through rapid decomposition due to high temperature. Several studies

in salt-affected soils proved that reclamation of alkali soils with amendment like gypsum application, and of saline soils by drainage installations will help in restoration of vegetation cover and increase of soil organic carbon. Association of grasses like *Leptochloa fusca* with multipurpose trees like *Prosopis juliflora* and *Acacia nilotica* in a unified agroforestry system in highly sodic soils in 4 to 7 year rotations has yielded biomass of about 24t ha<sup>-1</sup>. Our several experiments in different agro-eco regions further proved that carbon content in the soil continued to increase with age of plantation, type of tree and grass species and stocking rate.

Large scale plantation efforts on salt affected soils will be called up on wastelands using funding support from the climate change related projects of Govt. of India. This seems to one of the promising options to moderate global warming impact on agriculture in the near future. Large scale promotion of biofuel crops and multi-enter prize agriculture practices and rainwater harvesting and re-use will be called upon to moderate/mitigate climate change related risk in agriculture.

## References


1. Lal M (2001) Future climate change: Implications for Indian summer monsoon and its variability. *Current Science* 81(9): 1196-1207.
2. Yadav JSP, Singh K (1970) Tolerance of certain forest species to various degrees of salinity and alkali. *Indian Forester* 96(8): 587-599.
3. Singh G (2003) General review of opuntias in India. *Journal of the Professional association of the cactus*, pp. 30-46.
4. Singh G, Felker P (1998) Cactus now world foods. *Indian farming*, pp. 26-31.
5. Gill HS, Abrol IP, Samra JS (1987) *Forest ecology and management*. 22: 57-69
6. Samra JS, Singh G (2002) Drought management strategies. *Natural Resource Management Division, ICAR, New Delhi, India*, p. 65.
7. Singh G, Abrol IP, Cheema SS (1988) Agroforestry on alkali soils: effect of planting methods and amendments on initial growth, biomass accumulation and chemical composition of mesquite (*Prosopis juliflora*). *Agroforestry Systems* 7(2):135-160.



Creative Commons Attribution 4.0  
International License

For possible submissions Click Here

[Submit Article](#)



## Research in Medical & Engineering Sciences

### Benefits of Publishing with us

- High-level peer review and editorial services
- Freely accessible online immediately upon publication
- Authors retain the copyright to their work
- Licensing it under a Creative Commons license
- Visibility through different online platforms